


Regular Article

The role of perceived threats on mental health, social, and neurocognitive youth outcomes: A multicontextual, person-centered approach

May I. Conley¹ , Jasmine Hernandez¹, Joann M. Salvati², Dylan G. Gee¹ and Arielle Baskin-Sommers¹

¹Department of Psychology, Yale University, New Haven, CT, USA and ²Department of Psychiatry and Behavioral Sciences, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

Abstract

Perceived threat in youth's environments can elevate risk for mental health, social, and neurocognitive difficulties throughout the lifespan. However, few studies examine variability in youth's perceptions of threat across multiple contexts or evaluate outcomes across multiple domains, ultimately limiting our understanding of specific risks associated with perceived threats in different contexts. This study examined associations between perceived threat in youth's neighborhood, school, and family contexts at ages 9–10 and mental health, social, and neurocognitive outcomes at ages 11–12 within a large US cohort ($N = 5525$) enrolled in the Adolescent Brain Cognitive DevelopmentSM Study (ABCD Study[®]). Latent profile analysis revealed four distinct profiles: Low Threat in all contexts, Elevated Family Threat, Elevated Neighborhood Threat, and Elevated Threat in all contexts. Mixed-effect models and post hoc pairwise comparisons showed that youth in Elevated Threat profile had poorer mental health and social outcomes 2 years later. Youth in the Elevated Family Threat profile uniquely showed increased disruptive behavior symptoms, whereas youth in the Elevated Neighborhood Threat profile predominantly displayed increased sleep problems and worse neurocognitive outcomes 2 years later. Together, findings highlight the importance of considering perceptions of threat across multiple contexts to achieve a more nuanced developmental picture.

Keywords: mental health; neurocognition; perceived threat; social functioning; youth environments

(Received 13 June 2021; revised 13 December 2021; accepted 16 December 2021)

Introduction

Youth's perceptions of their environmental contexts, such as their neighborhoods, schools, and families, can play a critical role in academic, interpersonal, and occupational success (Brosschot et al., 2017; Burton et al., 1997; Caspi et al., 2012; Chen et al., 2004; Orstad et al., 2017; Repetti et al., 2002; Roosa et al., 2003, 2009; Thapa et al., 2013). Specifically, youth's perceptions of threat in their neighborhoods, schools, and families are associated with poorer mental health (e.g., Aldridge & McChesney, 2018; Aneshensel & Sucoff, 1996; Kasen et al., 1990), social functioning (e.g., Aldridge et al., 2018; Fite et al., 2010), and neurocognitive performance (e.g., Fay-Stammbach et al., 2014). In contrast to objective measures such as crime reports or census data, youth's perceptions of their neighborhoods, schools, and families have been found to be more predictive of mental health and peer relationship outcomes (Baranyi et al., 2021; Danese & Widom, 2020; El-Sheikh & Harger, 2001; Goldman-Mellor et al., 2016; Hadley-Ives et al., 2000). However, despite these general trends, several limitations in previous research on perceived threat constrain our understanding of the relationships between youth's

perceptions of threat in the neighborhood, school, and family and developmental outcomes. Notably, research often examines each context separately as if development occurs in only one setting; outcomes within a single domain (e.g., mental health, social, neurocognitive), hindering a more complete, cross-domain, understanding of the risks posed by perceived threats in each context; or homogenous samples limiting knowledge about the extent to which documented trends generalize across youth. Addressing these limitations is important in order to develop more holistic conceptualizations of the risks associated with perceived threat across these three primary contexts in which youth spend time (Hofferth & Sandberg, 2001).

Many foundational theories of development emphasize the importance of examining multiple contexts at different levels of proximity to youth (Bronfenbrenner, 1994; Bronfenbrenner & Morris, 2006; Cicchetti & Lynch, 1993; Lerner, 1991; Magnusson & Stattin, 1998). However, a large majority of research on the influence of perceived neighborhood, school, and family threats on development has examined each of these contexts separately. Yet, it is possible that youth have different experiences of threat within each context, such as perceiving threat in only one context versus more than one or not at all, which might impact development in unique ways (Cohodes et al., 2020; Youngblade et al., 2007). Moreover, youth environments grow increasingly complex throughout development, shifting from primarily family focused in early childhood to peer, school, and neighborhood

Corresponding author: May I. Conley, email: may.conley@yale.edu

Cite this article: Conley, M. I., et al. (2022). The role of perceived threats on mental health, social, and neurocognitive youth outcomes: A multicontextual, person-centered approach. *Development and Psychopathology*, 1–22, <https://doi.org/10.1017/S095457942100184X>

focused in later childhood and adolescence (Eccles & Roeser, 2009; Steinberg, 2005; Wigfield et al., 2006). Therefore, solely examining perceived threat in a single context, as was often done in previous research, may fail to capture precise relationships between perceived threats in youth contexts and developmental outcomes.

For example, perceptions of neighborhood and school threat independently have been associated with youth substance use and mental health (Kasen et al., 1990; Lambert et al., 2004; LaRusso et al., 2007; Wang & Degol, 2016). However, one study found that, when modeled together, perceived neighborhood threat, but not perceived school threat, was associated with adolescents' substance use and depressive symptoms (Nails et al., 2009). Another study showed that both perceived neighborhood and school threat relate to alcohol use (Friese et al., 2015), yet findings suggested there was an overall larger effect of perceived neighborhood threat on youth alcohol use. Although these two studies indicate that perceived neighborhood threat may be a specific risk for youth substance use, neither study examined an interaction between perceived neighborhood and school threat or considered the potential influence of perceived family threat, which also is associated with substance use outcomes (Repetti et al., 2002).

Additionally, outcomes associated with perceived neighborhood, school, and family threat span multiple domains of functioning (e.g., mental health, social, neurocognition). However, studies often examine a limited set of outcomes, often within a single domain, impeding comprehensive understanding of the influence of perceived threat in each context. For instance, perceived neighborhood threat has been associated with increased symptoms of depression, anxiety, and disruptive behavior disorders (DBD) (Aneshensel & Sucoff, 1996; Dawson et al., 2019), delinquency and proactive aggression (Byrnes et al., 2007; Fite et al., 2010), and poorer academic performance (Bowen et al., 2008; Williams et al., 2002) and verbal ability (Kohen et al., 2002). Yet, each of these studies only examined the few outcomes listed making it difficult to know whether the same youth who show symptoms of mental health difficulties also show decrements in social and neurocognitive functioning.

Consistent findings in the literature suggest that mental health, social, and neurocognitive functions are related, highlighting the importance of simultaneously examining a wide range of outcomes potentially related to perceived environmental threat. For example, a recent meta-analysis by Wagner et al. (2015) found that youth with depression perform worse on a range of neurocognitive tasks (e.g., sustained attention, working memory, verbal fluency) compared to youth without depression and neurocognitive deficits have been linked to interrupted social learning processes in DBD (Matthys et al., 2012). Other work has shown peer problems can be related to externalizing and internalizing in youth (Humphreys et al., 2013), and neurocognitive deficits may be associated with risk for peer problems (and vice versa; Holmes et al., 2016). Given these interrelations between mental health, social, and neurocognitive functions (Blanken et al., 2017; Klimes-Dougan & Garber, 2016; Ogilvie et al., 2011), it is possible that previous research examining the influences of perceived environmental threat has overlooked important associations by failing to evaluate multiple outcomes within the same sample.

Finally, although some work has examined perceived threat in multiple contexts or a wider range of outcomes, conclusions are often undermined by the use of homogenous samples. For example, one study conducted in a national sample of primarily White youth (81%) found that neither neighborhood nor school threats were associated with academic achievement (Youngblade et al., 2007).

This finding contrasts with another study conducted in a regionally restricted sample of primarily Black youth (88%) in Baltimore that showed both perceived neighborhood and school threats were associated with significantly worse reading and math achievement (Millam et al., 2010). Similarly, inconsistent patterns have been observed regarding mental health and behavioral outcomes. One study in a sample of predominantly Black youth in Alabama found that neighborhood threat attenuated the influence of family threat on internalizing and externalizing symptoms (Mrug & Windle, 2010). However, another study in racially/ethnically diverse youth (49.5% White, 26.3% Black, 14.3% Hispanic, and 9.8% other) in Texas found that neighborhood threat only attenuated the influence of family threat for internalizing, but not for externalizing, symptoms (Rosenfield et al., 2014). Together this work provides some evidence that unique combinations of perceived neighborhood, school, and family threat can differentially relate to unique outcomes during development (see Bacchini & Esposito, 2020 for review). However, it is not clear whether differences in findings across these studies should be attributed to combinations of perceived threat across multiple contexts, or whether differences in findings emerged from sample characteristics (i.e., racially or regionally homogenous samples). Therefore, before drilling down into comparisons between homogenous samples, research using a sociodemographically heterogeneous sample might be helpful for determining whether associations identified in previous work are representative of experiences more generally (Coley et al., 2018; Simmons et al., 2021).

Across decades of theoretical and empirical work, a robust literature has emerged that underscores associations between perceived environmental threat in different contexts and problems in mental health, social, and neurocognitive domains. Several mechanisms have been identified that purportedly link perceptions in different contexts to these outcomes. For example, neighborhoods, schools, and families can serve as important socialization contexts where youth learn rules and norms through processes, such as social modeling, observational learning, and interactions with family members, peers, and other community members (Bugental & Grusec, 2006). Youth's experiences, including their perceptions of threat, also can influence psychological and cognitive processes that result in various outcomes across multiple domains. For example, alterations in attentional (Pollak, 2015; Shackman & Pollak, 2014) and reward (Guyer et al., 2006) systems have been posited to underlie associations between youth's experiences of family threat and mental health and social functioning. In addition, biological models suggest youth's perceptions of threat get under the skin by influencing adaptations in nervous, endocrine, and immune systems (Danese & McEwen, 2012; Gunnar & Quevedo, 2007; Lupien et al., 2009; McEwen & Wingfield, 2003). Although research highlights several mechanisms at different levels of analysis that connect perceived threats with decrements in mental health, social, and neurocognitive development, little research has examined whether threat experienced across different contexts may differentially relate to outcomes across multiple domains in a sociodemographically diverse sample. It is therefore necessary to utilize a multicontextual, multioutcome approach to create a foundation for further investigation of how youth's perceptions of multicontextual threat interact with youth's changing biology to influence development.

As such, the goal of the present study was to examine heterogeneity in perceived threat in neighborhood, school, and family contexts in a large cohort of US youth. More specifically, we use the diverse Adolescent Brain Cognitive DevelopmentSM Study

(ABCD Study[®]) sample (Compton et al., 2019; Garavan et al., 2018) and longitudinal data. First, we implement a person-centered analysis (e.g., latent profile analysis [LPA]) to parse the heterogeneity across youth's perceptions of threat at baseline (ages 9–10; T1), rather than using a variable-centered approach that solely describes relationships related to a single variable (Howard & Hoffman, 2018; Magnusson & Stattin, 1998; Masyn, 2013; Vermunt & Magidson, 2002). Second, we use mixed-effect models and post hoc pairwise comparisons to investigate whether person-centered profiles of perceived threat predict mental health, social, and neurocognitive outcomes 2 years later (ages 11–12; T2). Most of the previous research examining the correlation between youth's perceptions of environmental threat and mental health, social functioning, and neurocognition is cross-sectional; therefore, using a longitudinal design advances our knowledge about the developmental sequelae of multicontextual perceived threat experiences in late childhood.

We hypothesized that profiles characterized by elevated perceived threat at baseline would be associated with decrements in developmental outcomes at the 2-year follow-up visit relative to profiles characterized by low perceptions of threat. Given that little progress has been made to identify person-centered profiles of perceived environmental threat, we did not have specific hypotheses about expected associations between the profiles and outcomes beyond the elevated threat versus low threat hypothesis described above. Rather, a key goal of this study was to explore whether heterogeneity in baseline perceived environmental threat differentially predicts mental health, social, and neurocognitive outcomes 2 years later and to create an empirical foundation for future research utilizing multicontextual and multioutcome approaches.

Materials and methods

Participants

Participants were children included in the ABCD Study Data Release 3.0 with complete baseline (T1; ages 9–10) and 2-year follow-up (T2; ages 11–12) data ($n = 6571$; 47.3% Female, 52.7% Male; 2.1% Asian; 11.9% Black; 19.3% Hispanic; 9.9% Other; 56.9% White; doi:10.15154/1519007; <https://nda.nih.gov/study.html?id=901>). A comparison based on baseline (T1) characteristics of participants with and without T2 data included in Data Release 3.0 is provided in Supplemental Table 1a–c. Participants were primarily recruited through schools in defined catchment areas for each of the 21 ABCD Study sites using a multistage probability sampling method to generate a sociodemographically diverse cohort (Garavan et al., 2018). Due to the geographic distribution of ABCD Study sites and potential self-selection bias in terms of enrollment, the sample is not perfectly representative of the US population overall (Compton, et al., 2019) and therefore may not perfectly generalize to all youth and families in the US.

The ABCD Study includes assessments of physical and mental health, neurocognition, biospecimens, substance use, culture and environment, and an extensive neuroimaging battery (Casey et al., 2018). All parents or caregivers provided written informed consent and children provided verbal assent for participation in the study (Clark et al., 2018). Baseline exclusionary criteria included a major neurological disorder (e.g., cerebral palsy, brain tumor, stroke, brain aneurysm, brain hemorrhage, subdural hematoma), multiple sclerosis, sickle cell disease, seizure disorders such as Lennox–Gastaut syndrome, Dravet syndrome, and Landau

Kleffner syndrome, a diagnosis of schizophrenia, moderate to severe autism spectrum disorder, intellectual disability, or history of substance use (i.e., all participants were substance use naive at baseline enrollment).

Assessments

ABCD Study data collection involves biennial visits with extensive (i.e., 6–7 h) mental health, social, and neurocognitive assessments and MRI scans, as well as brief (i.e., 2 h) yearly behavioral visits including interviews, questionnaires, and neurocognitive testing that touch on various domains, as well as biospecimen collection (Casey et al., 2018). Because only the biennial visits include an extensive evaluation across domains (Jernigan et al., 2018; <https://abcdstudy.org/scientists/protocols/>), in the present study we analyzed behavioral data collected from the baseline (T1) and 2-year (T2) follow-up visits across all 21 ABCD sites (doi:10.15154/1519007; <https://nda.nih.gov/study.html?id=901>).

Perceived environmental threat

Perceived environmental threat at T1 was estimated across three settings: neighborhood, school, and family. Correlations across perceived environmental threat and demographic variables are visualized in Supplemental Figure 1. Since the youth-report ABCD Neighborhood Safety/Crime Survey Modified from PhenX (NSC) (Echeverria et al., 2004; Mujahid et al., 2007) included only one item asking youth to indicate whether they strongly agreed or strongly disagreed with the statement: *My neighborhood is safe from crime*, perceived neighborhood threat was assessed using the mean of all youth- and parent-report items. The parent-report NSC included the exact same item described above and two additional items asking participants to indicate whether they strongly agreed or strongly disagreed with the statements: *I feel safe walking in my neighborhood, day or night* and *Violence is not a problem in my neighborhood*. All items were reverse-scored so that higher scores indicated *more* neighborhood threat (range = 1–5).

Perceived family threat was assessed using the youth-report summary score derived from the ABCD Youth Family Environment Scale-Family Conflict Subscale Modified from PhenX (FES-FCS) (Hoffman et al., 2019; Moos & Moos, 1994). The FES-FCS consisted of nine items evaluating the amount of conflict expressed by family members (e.g., *we fight a lot in our family; family members often criticize each other; family members sometimes hit each other*.) For each item, youth indicated whether each statement was true or false for most members of their family. All items were summed and scored with higher scores indicating *more* family conflict (range = 0–9).

Perceived school threat was assessed with the youth-report summary score derived from the School Environment subscale of the School Risk and Protective Factors (SRPF) Survey (Arthur et al., 2007). The SRPF School Environment subscale included six items evaluating youth's perceptions of the school climate related to safety and support (e.g., *I feel safe at my school; My teacher(s) notices when I am doing a good job and lets me know about it; Zucker et al., 2018*). For each item, youth indicated whether a statement was definitely true (YES! (4)) or definitely not true (NO! (1)). All items were summed and scored with higher scores indicating a *less* safe/supportive school environment (range = 1–24).

Mental health symptom outcomes

Mental health symptoms at T2 were evaluated using youth- and parent-reports. Different measures were available for a variety of mental health symptoms. Because some mental health symptoms may be difficult for parents and caregivers to detect (e.g., anxiety and depressive symptoms; Tandon et al., 2009), youth-report data were used where available. Supplemental analyses using only dimensional parent-report data were conducted and are available in Supplemental Materials.

Kiddie Schedule for Affective Disorders and Schizophrenia for DSM-5

The computerized Kiddie Schedule for Affective Disorders and Schizophrenia for DSM-5 (KSADS-5-PL) (Kaufman et al., 1997, 2013; Kobak et al., 2013) was used to assess symptoms associated with various mental health diagnoses. The KSADS-5-PL has high reliability and validity for assessing psychopathology in youth ages 6–18 and was optimized for use in ABCD Study data collection (Barch et al., 2018). Youth-report symptom counts were generated for anxiety disorders, conduct disorder, eating disorders, mood disorders, and suicidality. Parent-report symptom counts were generated for modules that were not included in the youth assessment including alcohol/substance use disorders, attention-deficit hyperactivity disorder (ADHD), DBD, homicidality, obsessive and compulsive disorders (OCD), other specified neurodevelopmental disorders, psychotic disorders, and trauma/stress disorders. Categorical diagnoses were included in supplemental analyses (see Supplemental Materials; Supplemental Table 5).

Achenbach System of Empirically Based Assessment Child Behavior Checklist

Broad dimensions of psychopathology symptoms (i.e., externalizing and internalizing *t*-scores) were evaluated using the Achenbach System of Empirically Based Assessment Child Behavior Checklist (CBCL). The CBCL is a well-established parent-report assessment used for identifying problem behavior in youth that is standardized and normed by age, sex, informant, and race/ethnicity (Achenbach, 2009). CBCL subscales were included in supplemental analyses (see Supplemental Materials).

Sleep Disturbance Scale for Children

Difficulties with sleep are a common feature of many mental health disorders, have been related to risk for substance use and subsequent psychopathology (Hasler et al., 2016), and have been related to perceived neighborhood and school threat (Meldrum et al., 2018). Total sleep problems were assessed using the ABCD Parent Sleep Disturbance Scale for Children (SDSC). The SDSC is a 26-item parent-report questionnaire that evaluates common sleep disorders in youth and has been validated for use in clinical and nonclinical samples (Bruni et al., 1996). The sum of the 26 items was used as a total score with higher scores indicating more difficulties with sleep.

Social outcomes

Social behavior and peer interactions were evaluated using measures from the ABCD culture and environment (Zucker et al., 2018) and mental health (Barch et al., 2018) assessments. All social outcomes were assessed at T2.

Peer network health protective scale

The degree of protection and support in peer networks was assessed with the ABCD Peer Network Health Protective Scale adapted from the Adolescent Social Network Assessment (Mason et al., 2004). This instrument prompted youth to indicate whether their closest friends had exhibited behaviors such as discouraging the use of substances, adopting healthy habits such as exercising or joining school clubs, or providing instrumental (e.g., school, money, transportation) and psychological support over the last 6 months. For items that were positively endorsed, youth selected how much encouragement their friends had provided from 1 to 10. Higher scores represented a greater protective peer network, and lower scores represented decreased peer network protection.

Youth peer behavioral profile

The Youth Peer Behavior Profile, derived from the Peer Behavior Profile/Peer Activities Questionnaire (Bingham et al., 1995), assessed two dimensions of youth peer networks: prosocial peers and rule breaking/delinquent peers. Each dimension corresponded to a 3-item subscale asking youth to report what proportion of their peers were involved in prosocial (e.g., excelling in school, playing sports) or rule breaking/delinquent (e.g., skipping school, shoplifting) behavior (range = 1, none or almost none–5, all or nearly all). Because the two subscales are not mutually exclusive, items for each subscale were summed separately.

ABCD peer experiences questionnaire

The ABCD Peer Experiences Questionnaire assessed 18 negative peer experiences such as overt, relational or reputational victimization from peers and perpetrating overt, relational or reputational aggression towards peers. For each item, youth indicated how often they had each experience in the past year using a five-point scale (range = 1, never–5, a few times a week). Total negative peer experiences were measured by summing all items with higher scores reflecting more negative peer experiences in the year prior to assessment.

Prosocial behavior scale

The ABCD Prosocial Behavior Scale from the Strengths and Difficulties Questionnaire (Goodman et al., 1998; Goodman & Scott, 1999) included three items that assessed youth's inclination to engage in behaviors that helped or supported others (e.g., being considerate of others' feelings). For each item, youth were asked to reflect on their behavior over the past 6 months and select whether each statement was Not True (0), Somewhat True (1), or Certainly True (2). All items were summed and higher scores indicated more prosocial behavior.

Neurocognitive outcomes

Neurocognitive outcomes were evaluated using behavioral data from 8 neurocognitive tasks administered at T2. All neurocognitive tasks were administered using iPads and have been previously detailed (Luciana et al., 2018).

The National Institutes of Health Toolbox Cognition Battery

Five cognitive tasks from the National Institutes of Health (NIH) Toolbox Cognition Battery were assessed at T2: a picture vocabulary task assessing language ability and vocabulary knowledge (Gershon et al., 2014), a Flanker task assessing attention and cognitive control (Fan et al., 2002), a picture sequence task assessing

episodic memory and visuospatial sequencing (Bauer et al., 2013; Dikmen et al., 2014), a pattern comparison task assessing visual information processing speed (Carlozzi et al., 2015, 2014, 2013), and an oral reading task assessing language and reading ability (Gershon et al., 2013). Because age-corrected scores are undergoing revision by the NIH Toolbox (Luciana et al., 2018), uncorrected standard scores were used to measure performance.

Rey Auditory Verbal Learning Test

The Rey Auditory Verbal Learning Test (RAVLT) was used to assess auditory verbal learning and memory (Lezak et al., 2004). The test involved five learning trials where participants were read a list of 15 unrelated words (list A). After each learning trial, participants were asked to recall as many words as possible. Next, participants were read a distractor list of 15 new words (list B) and were then asked to recall as many words as possible from the distractor list (list B). After the distractor trial, a recall trial was immediately assessed for words from the initial list (list A). After a 30-min delay where participants rest or complete nonverbal tasks, a final delayed recall trial is assessed for words from the initial list (list A). Here, we assessed performance (total correct) on the immediate and delayed recall trials (i.e., RAVLT Trials VI and VII).

Little Man Task

The Little Man Task (Acker & Acker, 1982) assesses visuospatial processing, perspective-taking and mental rotation. During administration of the task, participants saw a cartoon holding a briefcase in the left or right hand. The cartoon appeared in different presentations including right side up, upside down, facing the participant or facing away. Across these different presentations, the briefcase could be in either the left or right hand. Participants were instructed to indicate whether the briefcase was in the left or right hand. Performance was measured with percent accuracy (Luciana et al., 2018).

Social Influence Task

The Social Influence Task measures risk perception, propensity for risky decision making, and susceptibility to perceived peer influence (Knoll et al., 2015). Youth were presented with various risky scenarios (e.g., skiing really fast down a hill; hitchhiking; stealing honey out of a beehive) across 40 trials, and asked to rate each activity's risk using a slider ranging from "very low risk" to "very high risk." After initial ratings were submitted, youth were presented with a risk rating they were told was provided by a group of peers for the exact same scenarios. The peer rating condition was either 4 points lower (−4 condition), 2 points lower (−2 condition), 2 points higher (+2 condition), or 4 points higher (+4 condition) than the participant's initial rating. Participants were then asked to rate the riskiness of the scenario again. Social decision-making was measured for each of the four peer rating conditions by subtracting the mean initial rating from the mean final rating across all trials in each condition. Negative scores indicated more susceptibility to peer influence on the negative conditions (i.e., −4 and −2) and higher scores indicated more susceptibility to peer influence on the positive conditions (i.e., +4 and +2).

Analytic plan

All statistical analyses were performed in R version 3.6.3 (R Core Team, 2020). First, LPA was used to identify profiles of perceived environmental threat (i.e., family, school, neighborhood) at T1

using the tidyLPA package (Rosenberg et al., 2018), which utilizes the maximum likelihood estimator via the expectation–maximization algorithm (Scrucca et al., 2016). This analysis was performed on all participants who had complete perceived neighborhood, school, and family threat data ($n = 6530$). The optimal number of profiles was selected by comparing fit across six latent profile models (1–6 class models) and evaluating interpretability. The comparative fit of the models was assessed using the Bayesian Information Criterion (better fit indicated with smaller values; Schwarz, 1978), entropy (a measure of classification uncertainty ranging from 0–1 with more certainty and class discrimination indicated with values approaching 1 and values > 0.8 considered acceptable; Celeux & Soromenho, 1996; Weller et al., 2020), and bootstrapped likelihood ratio tests where a model k was considered to have preferable fit relative to a model $k-1$ when indicated with a significant p -value. Robustness of cluster results from the LPA was validated using k -means cluster analysis (see Supplemental Materials; Supplemental Figures 2–3). Participants were assigned to one profile for which their conditional probability of membership was the largest.

Next, associations between profile (a between-subjects factor) of T1 perceptions of threat and mental health, social, and neurocognitive outcomes at T2¹ were examined with mixed-effect models using the *lmer()* function from the *lme4* package (Bates et al., 2015). For all models, dependent variables and covariates were standardized and county-level crime and household income were included as fixed-effect covariates². Consistent with recommendations for the three-step model of LPA with multilevel data (Vermunt, 2010), ABCD Study site was included as a random intercept in all models to account for the multisite sampling design.

Finally, post hoc pairwise comparisons were performed using the *emmeans* package (Lenth, 2021) and used to evaluate differences in the magnitude of pairs of associations between outcome variables and separate latent profiles. Because the ABCD Study included siblings and twin pairs (Iacono et al., 2018), mixed-effect models and pairwise comparisons were performed using only one randomly selected child per family to avoid confounds introduced by family structure (final $n = 5525$; see Supplemental Table 2 for comparison of demographics for complete and incomplete datasets). Bonferroni correction was applied to account for the number of mixed-effect models generated for each outcome variable ($n = 33$). While visual inspection indicated that some outcome measures were not normally distributed (Supplemental Figures 4a–4c), mixed-effect models are robust violations of normality (Schielzeth et al., 2020; Verbeke & Molenberghs, 2013) and the adequacy of all models was evaluated using the *check_model()* function from the *performance* package (Lüdtke et al., 2021) to ensure that no modeling assumptions were violated.

Results

Person-centered profiles of perceived environmental threat at T1

Solutions for models with 1–6 latent classes were evaluated (Table 1). Bootstrapped likelihood ratio tests indicated that models with 2, 3, 4, and 5 classes showed improved fit relative to those with

¹Multiple T2 outcomes were not assessed at T1 (e.g., 4/5 T2 social outcomes not assessed at T1). However, where available, we evaluated whether membership in each profile at T1 predicted outcomes at T2 when accounting for T1 outcomes.

²Additional analyses included median family-income at the census tract level as a fixed-effect covariate to account for neighborhood-level socioeconomic resources.

Table 1. Model fit of the latent profile analysis

Classes	Log-likelihood	AIC	BIC	Entropy	BLRT	BLRT <i>p</i>
1	-27,795.51	55,603.01	55,643.72	1.00		
2	-27,010.86	54,041.72	54,109.56	0.82	1569.39	.01
3	-26,769.51	53,567.03	53,662.01	0.80	482.39	.01
4	-26,625.96	53,287.91	53,410.03	0.82	287.16	.01
5	-26,626.00	53,296.01	53,445.26	0.55	296.33	.01
6	-26,442.26	52,936.52	53,112.91	0.58	-0.21	.97

Note. AIC = Akaike information criterion; BIC = Bayesian information criterion; BLRT = bootstrapped likelihood ratio test. The selected class solution (4-class) is italicized and bolded.

one fewer class. Of these, the five-class solution was rejected because of low classification certainty as indicated by a low entropy value (0.55). Ultimately the four-class solution was selected as the best fitting model because it showed a preferable classification certainty (entropy = 0.82) and a lower BIC value (53,410.03) than the two and three-class solutions. As illustrated in Figure 1, the first profile was characterized by low threat ratings across all three contexts (profile 1 [Low Threat]; $n = 3953$; 71% of the sample) corresponding with low neighborhood threat (mean 1.80 ($SD = 0.58$)), low school threat (mean = 4.74 ($SD = 2.48$)), and low family threat (mean = 1.13 ($SD = 1.06$)). The second profile was characterized by elevated family threat (mean = 5.12 ($SD = 1.18$)), low neighborhood threat (mean = 2.04 ($SD = 0.75$)), and low school threat (mean = 5.42 ($SD = 2.51$)) (profile 2 [Elevated Family]; $n = 974$; 18% of the sample). The third profile was characterized by elevated neighborhood threat (mean = 3.75 ($SD = 0.47$)), low family threat (mean = 1.74 ($SD = 1.38$)), and low school threat (mean = 4.42 ($SD = 2.46$)) (profile 3 [Elevated Neighborhood]; $n = 458$; 8% of the sample). The fourth profile was characterized by elevated ratings across all three contexts (profile 4 [Elevated Threat]; $n = 140$; 2.5% of the sample) corresponding with elevated neighborhood threat (mean = 2.80 ($SD = 0.79$)), elevated school threat (mean = 12.72 ($SD = 2.15$)), and elevated family threat (mean = 3.91 ($SD = 1.79$)). Profile demographics are presented in Supplemental Table 3.

Profiles of perceived environmental threat predict developmental outcomes at T2

We examined whether each profile (at T1) differed on measures of mental health, social, and neurocognitive outcomes (at T2)^{3,4} using mixed-effect models and post hoc pairwise comparisons. Means and standard deviations for mental health, social, and neurocognitive data are presented in Supplemental Table 4. Results from the mixed-effect models and post hoc pairwise comparisons are detailed below (Table 2) and summarized in Figures 2–5.

Mental health symptom outcomes

Results within the mental health domain (Table 2 and Figure 2) demonstrated that membership in the Low Threat profile at T1

³Effects were consistent when controlling for T1 scores in linear mixed effect models. However, this was not possible for all models (e.g., 4/5 social outcomes) given changes between T1 and T2 protocols (<https://abcdstudy.org/scientists/protocols/>).

⁴Effects were consistent in additional analyses including neighborhood-level socioeconomic resources as a fixed-effect covariate along with county-level crime and household income.

was associated with lower ADHD, conduct, DBD, externalizing, internalizing, mood, neurodevelopmental, psychotic, suicidality, sleep problem, and trauma/stress symptoms at T2 relative to one or more of the elevated threat profiles. The Low Threat profile was not significantly different from the elevated threat profiles in alcohol/substance use, anxiety, eating, homicidality, or OCD symptoms at T2. Conversely, membership in the Elevated Threat profile at T1 was associated with increased ADHD, conduct, DBD, externalizing, internalizing, mood, sleep problem, suicidality, and trauma/stress symptoms at T2 relative to the Low Threat profile. Furthermore, membership in the Elevated Threat profile was related to even greater ADHD, conduct, externalizing, internalizing, sleep problem, suicidality, and trauma/stress symptoms relative to the Elevated Family profile and even greater ADHD, conduct, DBD, externalizing, internalizing, mood, and suicidality symptoms relative to the Elevated Neighborhood profile (summarized in Figure 3). Membership in the Elevated Family profile at T1 was significantly associated with increased conduct and DBD symptoms at T2 relative to the Elevated Neighborhood profile. Lastly, membership in the Elevated Neighborhood profile at T1 was significantly associated with increased externalizing and internalizing symptoms relative to the Low Threat profile, and increased sleep problems relative to both the Low Threat and Elevated Family profiles.

Social outcomes

Results within the social domain (Table 2 and Figure 4) demonstrated that membership in the Low Threat profile at T1 was associated with increased prosocial behavior and affiliation with prosocial peers and decreased negative peer experiences and affiliation with rule breaking peers relative to one or more of the elevated threat profiles at T2. By contrast, membership in the Elevated Threat profile at T1 was associated with increased negative peer experiences and affiliation with rule breaking peers and decreased prosocial behavior and affiliation with prosocial peers relative to all other profiles. Membership in the Elevated Family profile at T1 was associated with increased negative peer experiences and affiliation with rule breaking peers and decreased prosocial behavior relative to the Low Threat and Elevated Neighborhood profiles. Lastly, membership in the Elevated Neighborhood profile at T1 was not significantly associated with social behavior at T2.

Neurocognitive outcomes

Results within the neurocognitive domain (Table 2 and Figure 5) demonstrated that membership in the Low Threat profile at T1 was associated with increased performance on the Picture Memory, Picture Vocabulary, Reading Comprehension, and Visuospatial Processing tasks relative to one or more of the elevated profiles. Conversely, membership in the Elevated Threat profile was not significantly associated with neurocognitive performance at T2. Membership in the Elevated Family profile at T1 was associated with decreased performance on the Picture Memory, Picture Vocabulary, Reading Comprehension and Visuospatial Processing tasks relative to the Low Threat profile. Lastly, membership in the Elevated Neighborhood profile at T1 was associated with decreased performance on the Picture Vocabulary and Picture Memory tasks relative to all other profiles, and decreased performance on the Reading Comprehension task relative to the Low Threat profile.




Figure 1. Box plot showing classification results from the latent profile analysis. Bars reflect confidence intervals for profile centroids and boxes reflect the standard deviations within each profile.

Discussion

Heightened perceived threat in youth's environments is a well-known factor that can impact functioning. During development, youth's perceptions of threat may influence mental health, social, and neurocognitive outcomes. Using a large, diverse sample of US youth, the present study examined youth's perceptions of threat in their neighborhoods, schools, and families and evaluated whether profiles of perceived threat at ages 9–10 differentially predicted outcomes at ages 11–12. Broadly speaking, results from the profile analysis indicated variability in where and the extent to which youth perceive threat in their environments. Further, while the Low Threat profile was consistently related to generally better outcomes, differential effects were observed for specific outcomes when comparing profiles characterized by elevated perceived threat.

We identified four distinct profiles characterized by differences in perceived threat in the neighborhood, school, and family. As expected, there was a profile characterized by Low Threat and another profile characterized by Elevated Threat in all three contexts. The presence of two profiles that have fully aligned levels of perceived threat across all contexts is consistent with research showing that youth's experiences of stressors within and outside of the family can co-occur (Herrenkohl & Herrenkohl, 2007) and that neighborhood factors can be associated with family dynamics and school environments (see Minh et al., 2017 for review). The other two profiles were uniquely characterized by Elevated Family or Elevated Neighborhood threat, providing evidence that while some youth may perceive threat in multiple contexts, others have concentrated experiences of threat. These two profiles can be contextualized in light of other work showing variability in associations between family environments and neighborhoods (Furstenberg et al., 2000). For example, family threats can occur in the context of supportive neighborhoods (Silk et al., 2004) and many families in neighborhoods high in crime or other indicators of threat provide ample support for their children

(Beyers et al., 2003; Cuellar et al., 2015; Li & Fischer, 2017; Voisin et al., 2017). Together, results from the LPA identified discernible profiles of perceived threat in three primary contexts in which youth spend time (Hofferth & Sandberg, 2001) and suggest that the presence of threat in one context does not necessarily connote the presence of threat in another context or in all contexts.

The majority of the sample had the highest probability of membership in the Low Threat profile ($n = 3953$). Overall, this profile predicted fewer mental health symptoms and increased social functioning at ages 11–12. By contrast, membership in the smallest profile, Elevated Threat ($n = 140$), predicted increased mental health symptoms, negative peer experiences, affiliation with rule breaking peers, and decreased prosocial behavior and affiliation with prosocial peers. These findings are consistent with previous work showing that childhood adversity often co-occurs (Green et al., 2010) and that risk for long-term negative outcomes can increase with exposure to multiple stressors (Evans et al., 2013). Although the Low Threat versus Elevated Threat comparison is unsurprising, these findings dovetail with and potentially advance several prominent theories of early life adversity. Namely, cumulative risk approaches account for the number of adversities (including perceptions of threat) children experience (Dube et al., 2009; Evans et al., 2013), and propose that cumulative adverse experiences impact functioning via overload of youth's changing biological systems (Danese & McEwen, 2012). Yet, a common criticism of cumulative approaches is that they fall short in capturing differential effects of distinct types of adversity (Felitti et al., 1998). In contrast, dimensional (Sheridan & McLaughlin, 2014) or splitting approaches (Manly et al., 2001; Smith & Pollak, 2021; St Clair et al., 2015) propose that different aspects of youth's experiences may impact distinct outcomes via specific alterations in biological systems (Kuhlman et al., 2017; McLaughlin & Lambert, 2017; Palacios-Barrios & Hanson, 2019). Our findings potentially suggest that for some, but not all, youth, cumulative perceived threat experiences across contexts

Table 2. Results of mixed effect models and post hoc pairwise comparisons

	Mental health				
ADHD	β (SE)	CI	p	Pairwise comparisons	
Low Threat (LT)	-0.05 (0.02)	-0.09, -0.01	.028	LT vs. EF; LT vs. ET; EF vs. ET; EN vs. ET	
Elevated Family (EF)	0.18 (0.04)	0.11, 0.25	<.001		
Elevated Neighborhood (EN)	0.12 (0.05)	0.02, 0.23	.024		
Elevated Threat (ET)	0.39 (0.09)	0.22, 0.57	<.001		
Alcohol/substance use	β (SE)	CI	p	Pairwise comparisons	
Low Threat (LT)	-0.03 (0.02)	-0.06, 0.01	.141		
Elevated Family (EF)	0.11 (0.04)	0.03, 0.18	.007		
Elevated Neighborhood (EN)	0.06 (0.06)	-0.06, 0.17	.328		
Elevated Threat (ET)	0.15 (0.09)	-0.03, 0.34	.099		
Anxiety	β (SE)	CI	p	Pairwise comparisons	
Low Threat (LT)	-0.01 (0.02)	-0.05, 0.03	.562		
Elevated Family (EF)	0.07 (0.04)	-0.00, 0.14	.066		
Elevated Neighborhood (EN)	-0.01 (0.05)	-0.12, 0.10	.882		
Elevated Threat (ET)	0.04 (0.09)	-0.13, 0.22	.639		
Conduct	β (SE)	CI	p	Pairwise comparisons	
Low Threat (LT)	-0.06 (0.03)	-0.11, -0.01	.020	LT vs. EF; LT vs. ET; EF vs. EN; EF vs. ET; EN vs. ET	
Elevated Family (EF)	0.20 (0.04)	0.13, 0.26	<.001		
Elevated Neighborhood (EN)	0.07 (0.05)	-0.03, 0.17	.160		
Elevated Threat (ET)	0.58 (0.08)	0.42, 0.75	<.001		
DBD	β (SE)	CI	p	Pairwise comparisons	
Low Threat (LT)	-0.06 (0.02)	-0.11, -0.01	.010	LT vs. EF; LT vs. ET; EF vs. EN; EN vs. ET	
Elevated Family (EF)	0.25 (0.04)	0.18, 0.32	<.001		
Elevated Neighborhood (EN)	0.12 (0.05)	0.02, 0.23	.025		
Elevated Threat (ET)	0.42 (0.09)	0.24, 0.59	<.001		
Eating	β (SE)	CI	p	Pairwise comparisons	
Low Threat (LT)	-0.04 (0.02)	-0.08, 0.01	.095		
Elevated Family (EF)	0.07 (0.04)	0.00, 0.15	.039		
Elevated Neighborhood (EN)	0.08 (0.05)	-0.02, 0.18	.124		
Elevated Threat (ET)	0.22 (0.09)	0.05, 0.39	.010		
Homicidality	β (SE)	CI	p	Pairwise comparisons	
Low Threat (LT)	-0.01 (0.02)	-0.05, 0.02	.465		
Elevated Family (EF)	0.07 (0.04)	-0.00, 0.15	.050		
Elevated Neighborhood (EN)	0.02 (0.06)	-0.08, 0.13	.660		
Elevated Threat (ET)	0.01 (0.09)	-0.17, 0.19	.908		
Mood	β (SE)	CI	p	Pairwise comparisons	
Low Threat (LT)	-0.06 (0.02)	-0.11, -0.02	.008	LT vs. EF; LT vs. ET; EN vs. ET	
Elevated Family (EF)	0.20 (0.04)	0.13, 0.27	<.001		
Elevated Neighborhood (EN)	0.12 (0.05)	0.02, 0.22	.017		
Elevated Threat (ET)	0.35 (0.08)	0.18, 0.51	<.001		
Neurodevelopmental	β (SE)	CI	p	Pairwise comparisons	
Low Threat (LT)	-0.04 (0.02)	-0.09, 0.00	.056	LT vs. EF	
Elevated Family (EF)	0.16 (0.04)	0.09, 0.24	<.001		
Elevated Neighborhood (EN)	0.10 (0.05)	-0.00, 0.21	.061		
Elevated Threat (ET)	0.18 (0.09)	0.00, 0.35	.047		

(Continued)

Table 2 (Continued)

OCD	β (SE)	CI	<i>p</i>	Pairwise comparisons
Low Threat (LT)	−0.02 (0.02)	−0.06, 0.02	.294	
Elevated Family (EF)	0.08 (0.04)	0.00, 0.15	.038	
Elevated Neighborhood (EN)	0.07 (0.05)	−0.04, 0.17	.216	
Elevated Threat (ET)	0.11 (0.09)	−0.07, 0.28	.224	
Psychotic	β (SE)	CI	<i>p</i>	Pairwise comparisons
Low Threat (LT)	−0.03 (0.02)	−0.07, 0.01	.112	LT vs. EN
Elevated Family (EF)	0.06 (0.04)	−0.01, 0.14	.098	
Elevated Neighborhood (EN)	0.23 (0.05)	0.12, 0.34	<.001	
Elevated Threat (ET)	0.21 (0.09)	0.03, 0.38	.020	
Suicidality	β (SE)	CI	<i>p</i>	Pairwise comparisons
Low Threat (LT)	−0.03 (0.02)	−0.06, 0.01	.102	LT vs. ET; EF vs. ET; EN vs. ET
Elevated Family (EF)	0.11 (0.04)	0.03, 0.18	.004	
Elevated Neighborhood (EN)	−0.02 (0.06)	−0.12, 0.09	.760	
Elevated Threat (ET)	0.50 (0.09)	0.33, 0.68	<.001	
Trauma/Stress	β (SE)	CI	<i>p</i>	Pairwise comparisons
Low Threat (LT)	−0.03 (0.02)	−0.07, 0.01	.161	LT vs. ET; EF vs. ET
Elevated Family (EF)	0.07 (0.04)	−0.00, 0.15	.054	
Elevated Neighborhood (EN)	0.12 (0.05)	0.02, 0.23	.025	
Elevated Threat (ET)	0.28 (0.09)	0.11, 0.46	.001	
CBCL-externalizing	β (SE)	CI	<i>p</i>	Pairwise comparisons
Low Threat (LT)	−0.09 (0.03)	−0.14, −0.04	.001	LT vs. EF; LT vs. EN; LT vs. ET; EF vs. ET; EN vs. ET
Elevated Family (EF)	0.25 (0.04)	0.18, 0.32	<.001	
Elevated Neighborhood (EN)	0.19 (0.05)	0.08, 0.29	<.001	
Elevated Threat (ET)	0.55 (0.09)	0.38, 0.72	<.001	
CBCL-internalizing	β (SE)	CI	<i>p</i>	Pairwise comparisons
Low Threat (LT)	−0.05 (0.03)	−0.11, 0.02	.172	LT vs. EN; LT vs. ET; EF vs. ET; EN vs. ET
Elevated Family (EF)	0.07 (0.04)	−0.01, 0.14	.071	
Elevated Neighborhood (EN)	0.18 (0.05)	0.07, 0.28	.001	
Elevated Threat (ET)	0.51 (0.09)	0.33, 0.68	<.001	
Sleep problems	β (SE)	CI	<i>p</i>	Pairwise comparisons
Low Threat (LT)	−0.04 (0.04)	−0.11, 0.04	.328	LT vs. EN; LT vs. ET; EF vs. EN; EF vs. ET
Elevated Family (EF)	0.11 (0.04)	0.04, 0.18	.003	
Elevated Neighborhood (EN)	0.25 (0.05)	0.15, 0.36	<.001	
Elevated Threat (ET)	0.30 (0.09)	0.13, 0.47	<.001	
Social				
Negative peer experiences	β (SE)	CI	<i>p</i>	Pairwise comparisons
Low Threat (LT)	−0.08 (0.02)	−0.12, −0.03	.002	LT vs. EF; LT vs. EN; EF vs. EN; EF vs. ET; EN vs. ET
Elevated Family (EF)	0.25 (0.04)	0.18, 0.33	<.001	
Elevated Neighborhood (EN)	0.14 (0.05)	0.03, 0.24	.011	
Elevated Threat (ET)	0.57 (0.09)	0.39, 0.74	<.001	
Prosocial behavior (Youth)	β (SE)	CI	<i>p</i>	Pairwise comparisons
Low Threat (LT)	0.04 (0.03)	−0.01, 0.09	.085	LT vs. EF; LT vs. ET; EF vs. EN; EF vs. ET; EN vs. ET
Elevated Family (EF)	−0.21 (0.04)	−0.29, −0.14	<.001	
Elevated Neighborhood (EN)	0.03 (0.05)	−0.08, 0.14	.577	
Elevated Threat (ET)	−0.58 (0.09)	−0.75, −0.40	<.001	

(Continued)

Table 2 (Continued)

Prosocial peers	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.01 (0.04)	-0.07, 0.09	.845	LT vs. ET; EF vs. ET; EN vs. ET
Elevated Family (EF)	-0.04 (0.04)	-0.12, 0.03	.232	
Elevated Neighborhood (EN)	0.03 (0.05)	-0.08, 0.13	.636	
Elevated Threat (ET)	-0.43 (0.09)	-0.60, -0.26	<.001	
Protective peer network	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	-0.01 (0.03)	-0.06, 0.04	.664	
Elevated Family (EF)	-0.03 (0.04)	-0.11, 0.04	.400	
Elevated Neighborhood (EN)	0.09 (0.05)	-0.01, 0.20	.081	
Elevated Threat (ET)	-0.10 (0.09)	-0.27, 0.08	.278	
Rule breaking peers	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	-0.06 (0.03)	-0.12, 0.01	.093	LT vs. EF; LT vs. ET; EF vs. ET; EN vs. ET
Elevated Family (EF)	0.19 (0.04)	0.12, 0.26	<.001	
Elevated Neighborhood (EN)	0.09 (0.05)	-0.01, 0.20	.086	
Elevated Threat (ET)	0.39 (0.09)	0.21, 0.56	<.001	
Neurocognitive				
Flanker	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.04 (0.04)	-0.03, 0.11	.233	
Elevated Family (EF)	-0.12 (0.04)	-0.19, -0.04	.002	
Elevated Neighborhood (EN)	-0.11 (0.05)	-0.21, 0.00	.051	
Elevated Threat (ET)	-0.18 (0.09)	-0.35, -0.00	.045	
Pattern comparison	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.05 (0.04)	-0.02, 0.12	.165	
Elevated Family (EF)	-0.11 (0.04)	-0.18, -0.03	.005	
Elevated Neighborhood (EN)	-0.16 (0.05)	-0.27, -0.05	.003	
Elevated Threat (ET)	-0.23 (0.09)	-0.40, -0.05	.011	
Picture memory	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.06 (0.03)	0.01, 0.12	.028	LT vs. EF; LT vs. EN; EF vs. EN
Elevated Family (EF)	-0.12 (0.04)	-0.19, -0.05	.001	
Elevated Neighborhood (EN)	-0.28 (0.05)	-0.38, -0.17	<.001	
Elevated Threat (ET)	-0.19 (0.09)	-0.36, -0.01	.033	
Picture vocabulary	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.09 (0.04)	0.02, 0.17	.014	LT vs. EF; LT vs. EN; EF vs. EN; EN vs. ET
Elevated Family (EF)	-0.15 (0.03)	-0.22, -0.08	<.001	
Elevated Neighborhood (EN)	-0.31 (0.05)	-0.41, -0.22	<.001	
Elevated Threat (ET)	-0.07 (0.08)	-0.23, 0.10	.438	
Reading comprehension	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.07 (0.03)	0.01, 0.12	.014	LT vs. EF; LT vs. EN
Elevated Family (EF)	-0.13 (0.04)	-0.20, -0.05	<.001	
Elevated Neighborhood (EN)	-0.19 (0.05)	-0.29, -0.08	<.001	
Elevated Threat (ET)	-0.14 (0.09)	-0.31, 0.03	.111	
Visuospatial processing	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.05 (0.03)	-0.00, 0.10	.067	LT vs. EF
Elevated Family (EF)	-0.12 (0.04)	-0.19, -0.05	.001	
Elevated Neighborhood (EN)	-0.08 (0.05)	-0.19, 0.02	.115	
Elevated Threat (ET)	-0.26 (0.09)	-0.43, -0.09	.003	

(Continued)

Table 2 (Continued)

Verbal recall (immediate)	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.03 (0.03)	−0.02, 0.09	.232	
Elevated Family (EF)	−0.04 (0.04)	−0.11, 0.03	.296	
Elevated Neighborhood (EN)	−0.16 (0.05)	−0.27, −0.06	.003	
Elevated Threat (ET)	−0.07 (0.09)	−0.24, 0.11	.454	
Verbal recall (delay)	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.03 (0.03)	−0.03, 0.08	.327	
Elevated Family (EF)	−0.03 (0.04)	−0.10, 0.05	.447	
Elevated Neighborhood (EN)	−0.11 (0.05)	−0.22, −0.00	.046	
Elevated Threat (ET)	−0.03 (0.09)	−0.21, 0.14	.723	
Social decision making (−4 condition)	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.03 (0.03)	−0.03, 0.09	.297	
Elevated Family (EF)	−0.04 (0.04)	−0.12, 0.03	.242	
Elevated Neighborhood (EN)	−0.09 (0.05)	−0.20, 0.01	.084	
Elevated Threat (ET)	0.16 (0.09)	−0.02, 0.33	.074	
Social decision making (−2 condition)	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	0.04 (0.02)	−0.01, 0.08	.135	
Elevated Family (EF)	−0.04 (0.04)	−0.11, 0.03	.278	
Elevated Neighborhood (EN)	−0.16 (0.05)	−0.26, −0.05	.004	
Elevated Threat (ET)	0.07 (0.09)	−0.11, 0.24	.445	
Social decision making (+2 condition)	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	−.02 (.02)	−.06, .03	.443	
Elevated Family (EF)	.03 (.04)	−.05, .10	.494	
Elevated Neighborhood (EN)	.00 (.05)	−.10, .11	.932	
Elevated Threat (ET)	−.15 (.09)	−.33, .02	.086	
Social decision making (+4 condition)	β (SE)	CI	p	Pairwise comparisons
Low Threat (LT)	−.03 (.03)	−.08, .02	.300	
Elevated Family (EF)	.04 (.04)	−.03, .12	.252	
Elevated Neighborhood (EN)	.07 (.06)	−.04, .18	.195	
Elevated Threat (ET)	−.15 (.09)	−.33, .03	.095	

Note. All models include household income and county-level crime as covariates. Bold p values indicate significance following Bonferroni correction. Listed pairwise comparisons (far right column) indicate significant differences ($p < .05$) between profiles. ADHD = attention deficit hyperactivity disorder; CBCL = Child Behavior Checklist; DBD = disruptive behavior disorders; OCD = obsessive and compulsive disorders.

may explain outcomes more than perceived threat in any single context alone and are associated with the largest mental health and social risks. Better understanding the ways in which diffuse (e.g., multiple contexts) versus concentrated (i.e., one context) perceptions of threat confer risk for mental health and social difficulties will allow for greater accuracy and precision in detecting risk and targeting intervention efforts. At the same time, the Elevated Threat profile did not show significantly increased risk for neurocognitive decrements relative to the Elevated Family or Elevated Neighborhood profiles. Accordingly, different aspects of perceived threat experiences, such as concentrated threat in one context, may be more related to neurocognitive outcomes than cumulative experiences alone.

The second largest profile, Elevated Family ($n = 974$) captured nearly a fifth of the sample. The Elevated Family profile predicted increased mental health symptoms and decreased neurocognitive

and social functioning relative to the Low Threat profile. Additionally, the Elevated Family profile showed increased conduct (youth-report) and DBD (parent-report) symptoms and decreased social functioning relative to the Elevated Neighborhood profile. The prominence of the family in predicting disruptive behavior symptoms is consistent with previous work detailing family processes characterized by harsh, coercive, or psychologically-controlling parenting as risks for disruptive and aggressive behavior (Gard et al., 2017; Kawabata et al., 2011; Latham et al., 2017; Oliver, 2015; Patterson, 1982; Pinquart, 2017; Romano et al., 2005). Childhood externalizing disorders, which have been linked with social impairment (Greene et al., 2002) and persistent antisocial and criminal behavior throughout the lifespan (Copeland et al., 2017; Fergusson et al., 2005; Herba et al., 2007; Rivenbark et al., 2018), can increase risk for other forms of psychopathology including depression and anxiety, and




Figure 2. Bar plot showing mental health symptoms at T2 as a function of each perceived environmental threat profile at T1. ADHD = attention deficit hyperactivity disorder; CBCL = Child Behavior Checklist; DBD = disruptive behavior disorders.

substance abuse (Copeland et al., 2017; Herba et al., 2007), posing high cost for the individual, their community, and society-at-large.

Family socialization models propose that the family provides a context where children learn a good deal about intra- and interpersonal functioning through social modeling and observational learning during their interactions with caregivers and the larger family climate (Denham et al., 1997; Morris et al., 2007; Parke, 1994). Threat in the family context may contribute to aberrations in self-regulation (e.g., increased impulsivity), which is implicated in the onset and maintenance of externalizing symptoms (Cappadocia et al., 2009; Fairchild et al., 2019; Masi et al., 2014). Therefore, known perceptions of family threat may warrant specialized screening for externalizing disorders, increased targeted

social skills development, and self-regulation interventions, and more general parenting skills and family support interventions.

Lastly, the Elevated Neighborhood profile ($n = 458$) was characterized by increased mental health symptoms relative to the Low Threat profile and increased sleep problems relative to the Low Threat and Elevated Family profiles. In addition, membership in the Elevated Neighborhood profile also was associated with the largest decreases in picture memory (i.e., episodic memory) and picture vocabulary (i.e., language ability and vocabulary knowledge) performance relative to all other profiles. These results are consistent with previous work showing that youth who perceive less neighborhood safety are at risk for insufficient sleep (Bagley et al., 2016; Mayne et al., 2021; Meldrum et al., 2018), which is




Figure 3. Schematic depicting largest unique effects across elevated profiles. Outcomes displayed reflect associations that were significantly greater in magnitude than at least one of the other elevated profiles and were not significantly lower than any of the other elevated profiles. Outcomes displayed for more than one profile (i.e., DBD, sleep problems) were not significantly different between those profiles. Icons in red represent elevated perceived threat in that context. ADHD = attention deficit hyperactivity disorder; DBD = disruptive behavior disorders.

theorized to be one factor linking neighborhood conditions to disparities in multiple physical and mental health outcomes (Hale et al., 2013; Jackson et al., 2015; Roberts & Duong, 2017; Shan et al., 2015). Notably, poor sleep quality is one factor known to impair neurocognitive functioning (Arnsten et al., 2015; Telzer et al., 2013).

Youth in the Elevated Neighborhood profile showed the largest decreases in neurocognitive performance. This finding aligns with documented associations between exposure to neighborhood violence and decrements in performance on vocabulary and reading tasks (Sampson et al., 2008; Sharkey, 2010). It is possible that in the face of perceived neighborhood threat it may benefit youth to prioritize other cognitive processes such as attention to negative stimuli in the environment (McCoy et al., 2016). Although this might support short-term adaptation to a specific, stressful environment (Amso, 2020), over time negative attentional bias may contribute to depression (Gotlib & Joormann, 2010) or other mental health risks (Mogg & Bradley, 2005). As cognitive performance, especially language ability, is a strong predictor of academic (Young et al., 2002) and vocational outcomes (Maughan, 1995), it may be advisable to assess school performance in youth experiencing neighborhood threat and increase academic support and skills to optimize learning. Additionally, youth who report experiencing threat in their neighborhoods may benefit from other skills and strategies to cope with stress, manage sleep, and increase their feelings of safety (Rasmussen et al., 2004). While we did not observe significantly increased mental health risk in the Elevated Neighborhood profile relative to the two other elevated profiles, these youth do show risk for mental health difficulties relative to

the Low Threat profile (i.e., increased internalizing and externalizing symptoms) and future work is needed to elucidate complex links between neighborhood factors, neurocognition, and physical and mental health outcomes.

Before concluding, several limitations and considerations for future research should be noted. First, while we explore how experiences at ages 9–10 predict outcomes at ages 11–12, the correlational nature of the analysis cannot speak directly to potential shared antecedents, mechanisms, or moderators that may influence the observed associations. For example, there may be shared antecedent factors (e.g., genetics, history of stressful life events, and so forth) that make individuals more susceptible to perceiving threats and showing decrements in mental health, social functioning, or neurocognitive performance (Cicchetti, 2010; Germine et al., 2016; Harkness et al., 2006; Wade et al., 2019). Additionally, several mechanisms at multiple levels of analysis (e.g., social-contextual, cognitive, neurobiological, and so forth) have been identified as important for linking youth’s experiences of threat to functioning across mental health, social, and neurocognitive domains (Bugental & Grusec, 2006; Danese & McEwen, 2012; Gunnar & Quevedo, 2007; Guyer et al., 2006; Lupien et al., 2009; Pollak, 2015; Shackman & Pollak, 2014). Future work is needed to investigate potential mechanistic pathways by which differential associations emerge between multicontextual perceived threats and specific developmental outcomes. Further, the present study does not speak to potential moderating factors that may attenuate or increase vulnerability for certain outcomes given perceptions of threat. For instance, supportive, close relationships (Colich et al., 2021; Rudolph et al., 2020), participation in




Figure 4. Bar plot showing social outcomes at T2 as a function of each perceived environmental threat profile at T1.

community organizations (Garmezy, 1991), and individual-level characteristics such as high self-esteem or achievement motivation can serve as protective factors for youth facing hardship (Hostinar & Miller, 2019; Masten, 2001; Masten & Narayan, 2012). A better understanding of protective and vulnerability factors that may influence complex associations between youth's perceptions of multicontextual threat and developmental outcomes will be important for optimizing intervention efforts to promote resilience.

Second, youth psychopathology and normative variation in certain symptoms (e.g., anxiety) can influence youth's perceptions of threat across multiple contexts (Puliafico & Kendall, 2006) and bidirectional links between youth behavior and different contexts, especially the family, are well-documented (Burt

et al., 2005; Masten & Cicchetti, 2010; Wiggins et al., 2015). Given limitations associated with lagged designs examining within-person variance with 3 or fewer waves of data (Orth et al., 2021; Usami et al., 2019), longitudinal evaluation of directionality between youth's perceptions of threat and mental health and behavior was not possible in the present study due to variation in which measures are assessed across ABCD Study visits for which data are currently available (<https://abcdstudy.org/scientists/protocols/>). However, research with future waves of ABCD Study data can be used to better understand the associations between perceived threat and outcomes at later stages of development and the directionality between youth's perceptions of threats in different contexts and youth mental health and behavioral outcomes.




Figure 5. Bar plot showing neurocognitive outcomes at T2 as a function of each perceived environmental threat profile at T1.

Third, although perceptions of threat in the neighborhood, school, and family have been shown to be distinct risk factors and influences on development (Danese & Widom, 2020; El-Sheikh & Harger, 2001; Goldman-Mellor et al., 2016; Hadley-Ives et al., 2000), there are contexts and indicators of threat that were not assessed in the ABCD Study. For example, youth become increasingly peer-oriented throughout development (Eccles & Roeser, 2009; Steinberg, 2005; Wigfield et al., 2006) and future research including measures of perceived peer threats will be important for further understanding youth's experiences of threat throughout development. In addition, the present study was unable to account for multiple objective indicators of threat. The ABCD Study battery does not assess objective school or family threats (e.g., objective reports of school violence or childhood

maltreatment) and residential history-derived crime-report data, which was included as a covariate in our analyses, is only available at the county-level at this time. Future releases of ABCD Study data may have more fine-grained objective measures (i.e., crime-report data at the neighborhood level) and future work comparing the influence of youth's perceptions to objective indicators of threat is needed.

Fourth, while we used a conservative threshold for evaluating significance, the overall magnitude of observed effects is small using traditional heuristics (Cohen, 1988). That said, recent work using ABCD Study data has suggested new benchmarks for effect size such as small $\leq .05$, medium = .06–.15, large = .16–.25 (Owens et al., 2021). Nonetheless, replication is needed to evaluate the reliability of associations between membership in the perceived threat

profiles and outcomes. Although it is likely that small, yet meaningful, environmental and biological factors work in concert to influence development (Dick et al., 2021), some of the smaller effects observed in the present study may be related to variability in the sequelae of perceived environmental threat. Responses to perceived threat are theorized to be informed by interactions between demands (i.e., perceptions of uncertainty, danger, and effort) and resources (i.e., coping skills/ability, dispositional factors, social support) (Jamieson et al., 2018). In the present study, youth who perceived elevated environmental threat (i.e., demands) and had adequate coping skills/ability, dispositional factors, or social/other support (i.e., resources) may not have shown the same mental health, social, or neurocognitive outcomes as youth who perceived the exact same profile of environmental threat without adequate coping and support resources. Further research exploring individual differences in which youth are most affected by perceived environmental threat will be important for increased understanding of the developmental sequelae of perceived environmental threat. Moreover, clinicians and other professionals who work with youth should consider the availability of resources that may interact with youth's perceptions of environmental threat when developing interventions.

Lastly, while the use of a large sample that estimates the diversity of the US on race and ethnicity, socioeconomic status, and urbanicity is a strength of the current study, the ABCD Study sample is limited with regard to the representation of rural families and is not perfectly representative of the US population overall (Compton et al., 2019). Moreover, participants included in the current study (i.e., those with T2 data included in Data Release 3.0) were significantly older and had a higher proportion of youth who identified as White, and parents who indicated higher levels of education and household income relative to the rest of the ABCD Study sample (Supplemental Table 1a–c). Further, youth from different backgrounds may have different experiences of threat across multiple contexts and may perceive threat in their environments for different reasons. In the present study, the Elevated Neighborhood profile had a significantly higher proportion of youth who identified as Black or Hispanic, and whose parents reported lower education levels and household income relative to the other three profiles. This pattern is consistent with other work showing that people of color are overrepresented in lower socioeconomic status neighborhoods (Williams & Collins, 2001) and encounter the highest rates of exposure to violence (Friedson & Sharkey, 2015; McNulty & Bellair, 2003; Williams & Jackson, 2005). While exposure to violence may be one factor that influences youth's perceptions of neighborhood threat, other research shows that youth of color can have disproportionate contact with the criminal legal system relative to White youth despite similar or lower levels of criminal offending (Padgaonkar et al., 2021), which also may influence perceptions of neighborhood threat. That said, there was sociodemographic variability across all four profiles suggesting that sociodemographic factors do not solely shape youth's perceptions of threat across different contexts. Ultimately, more research is needed to understand how youth's intersectional identities (Cole, 2009; Crenshaw, 1993) interact with their experiences in different contexts to influence perceptions of threat and confer risk for or resilience against difficulties across multiple domains.

Increased knowledge of heterogeneity in youth experiences of perceived environmental threat is important for moving closer to the goal of more fully understanding multifaceted associations between youth environments and developmental outcomes.

Understanding youth's specific experiences of threat in different contexts is important for tailoring skills and strategies to focus on the individual to help support youth in coping and meeting developmental goals. Here our findings implicate perceived threat in youth's environments as a common risk factor that cuts across diagnostic boundaries and domains of functioning. Further, specific profiles of threat may pose greater risk for certain types of outcomes. Together, results underscore calls to action for clinicians to not only treat individuals, but also advocate for research-based policy that bolsters safe and supportive environments that promote positive youth development.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S095457942100184X>

Data availability statement. Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development (ABCD) Study (<https://abcdstudy.org>), held in the NIMH Data Archive (NDA). This is a multi-site, longitudinal study designed to recruit more than 10,000 children age 9–10 and follow them over 10 years into early adulthood.

Funding statement. The ABCD Study is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at <https://abcdstudy.org/federal-partners.html>. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/consortium_members/. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators. The ABCD data repository grows and changes over time. The ABCD data used in this report came from DOI: 10.15154/1519007.

Conflicts of interest. None.

References

- Achenbach, T. M. (2009). *The Achenbach System of Empirically Based Assessment (ASEBA): Development, findings, theory, and applications*. University of Vermont, Research Center for Children, Youth, & Families.
- Acker, W., & Acker, C. (1982). *Bexley Maudsley automated processing screening and Bexley Maudsley category sorting test manual*. NFER-Nelson, Windsor, England.
- Aldridge, J. M., & McChesney, K. (2018). The relationships between school climate and adolescent mental health and wellbeing: A systematic literature review. *International Journal of Educational Research*, 88, 121–145. <https://doi.org/10.1016/j.ijer.2018.01.012>
- Aldridge, J. M., McChesney, K., & Afari, E. (2018). Relationships between school climate, bullying and delinquent behaviours. *Learning Environments Research*, 21(2), 153–172. <https://doi.org/10.1007/s10984-017-9249-6>
- Amso, D. (2020). Neighborhood poverty and brain development: Adaptation or maturation, fixed or reversible? *JAMA Network Open*, 3(11), e2024139. <https://doi.org/10.1001/jamanetworkopen.2020.24139>
- Aneshensel, C. S., & Sucoff, C. A. (1996). The neighborhood context of adolescent mental health. *Journal of Health and Social Behavior*, 37(4), 293–310.
- Arnsten, A. F. T., Raskind, M. A., Taylor, F. B., & Connor, D. F. (2015). The effects of stress exposure on prefrontal cortex: Translating basic research into successful treatments for post-traumatic stress disorder. *Neurobiology of Stress*, 1, 89–99. <https://doi.org/10.1016/j.yjnstr.2014.10.002>

- Arthur, M. W., Briney, J. S., Hawkins, J. D., Abbott, R. D., Brooke-Weiss, B. L., & Catalano, R. F. (2007). Measuring risk and protection in communities using the Communities That Care Youth Survey. *Evaluation and Program Planning*, 30(2), 197–211. <https://doi.org/10.1016/j.evalprogplan.2007.01.009>
- Bacchini, D., & Esposito, C. (2020). Growing up in violent contexts: Differential effects of community, family, and school violence on child adjustment. In Balvin N., & Christie D., Children and peace. Peace psychology book series, Springer Cham, https://doi.org/10.1007/978-3-030-22176-8_10
- Bagley, E. J., Tu, K. M., Buckhalt, J. A., & El-Sheikh, M. (2016). Community violence concerns and adolescent sleep. *Sleep Health*, 2(1), 57–62. <https://doi.org/10.1016/j.sleh.2015.12.006>
- Baranyi, G., Di Marco, M. H., Russ, T. C., Dibben, C., & Pearce, J. (2021). The impact of neighbourhood crime on mental health: A systematic review and meta-analysis. *Social Science & Medicine*, 282, 114106. <https://doi.org/10.1016/j.socscimed.2021.114106>
- Barch, D. M., Albaugh, M. D., Avenevoli, S., Chang, L., Clark, D. B., Glantz, M. D. . . . Sher, K. J. (2018). Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Developmental Cognitive Neuroscience*, 32, 55–66. <https://doi.org/10.1016/j.dcn.2017.10.010>
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software, Articles*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Bauer, P. J., Dikmen, S. S., Heaton, R. K., Mungas, D., Slotkin, J., & Beaumont, J. L. (2013). III. NIH Toolbox cognition battery (CB): Measuring episodic memory. *Monographs of the Society for Research in Child Development*, 78(4), 34–48. <https://doi.org/10.1111/mono.12033>
- Bingham, C. R., Fitzgerald, H. E., & Zucker, R. A. (1995). *Peer Behavior Profile/Peer Activities Questionnaire*. Unpublished questionnaire. Department of Psychology, Michigan State University, East Lansing.
- Beyers, J. M., Bates, J. E., Pettit, G. S., & Dodge, K. A. (2003). Neighborhood structure, parenting processes, and the development of youth's externalizing behaviors: A multilevel analysis. *American Journal of Community Psychology*, 31(1–2), 35–53. <https://doi.org/10.1023/a:1023018502759>
- Blanken, L. M. E., White, T., Mous, S. E., Basten, M., Muetzel, R. L., Jaddoe, V. W. V. . . . Tiemeier, H. (2017). Cognitive functioning in children with internalising, externalising and dysregulation problems: A population-based study. *European Child & Adolescent Psychiatry*, 26(4), 445–456. <https://doi.org/10.1007/s00787-016-0903-9>
- Bowen, G. L., Rose, R. A., Powers, J. D., & Glennie, E. J. (2008). The joint effects of neighborhoods, schools, peers, and families on changes in the school success of middle school students. *Family Relations*, 57(4), 504–516. <https://doi.org/10.1111/j.1741-3729.2008.00518.x>
- Bronfenbrenner, U. (1994). Ecological models of human development. In Gauvain M., & Cole M. (Eds.), *International encyclopedia of education*. vol. 3, 2nd. Elsevier Oxford: 37–43.
- Bronfenbrenner, U., & Morris, P. A. (2006). The bioecological model of human development. In Lerner R. M., & Damon W. (Eds.), *Handbook of child psychology: Theoretical models of human development*. John Wiley & Sons Inc: 793–828.
- Brosschot, J. F., Verkuil, B., & Thayer, J. F. (2017). Exposed to events that never happen: Generalized unsafety, the default stress response, and prolonged autonomic activity. *Neuroscience and Biobehavioral Reviews*, 74(Pt B), 287–296. <https://doi.org/10.1016/j.neubiorev.2016.07.019>
- Bruni, O., Ottaviano, S., Guidetti, V., Romoli, M., Innocenzi, M., Cortesi, F. . . . Giannotti, F. (1996). The Sleep Disturbance Scale for Children (SDSC). Construction and validation of an instrument to evaluate sleep disturbances in childhood and adolescence. *Journal of Sleep Research*, 5(4), 251–261. <https://doi.org/10.1111/j.1365-2869.1996.00251.x>
- Bugental, D. B., & Grusec, J. E. (2006). Socialization processes. In Damon W., Lerner R. M., & Eisenberg N. (Eds.), *Handbook of child psychology*. vol. 3, John Wiley & Sons Inc: 366–428.
- Burt, S. A., McGue, M. A. T. T., Krueger, R. F., & Iacono, W. G. (2005). How are parent-child conflict and childhood externalizing symptoms related over time? Results from a genetically informative cross-lagged study. *Development and Psychopathology*, 17(1), 145. <https://doi.org/10.1017/S095457940505008X>
- Burton, L. M., Price-Spratlen, T., & Spenser, M. B. (1997). On ways of thinking about measuring neighborhoods: Implications for studying context and development outcomes for children. In Brooks-Gunn, J., Duncan, G., & Lawrence Aber, J. (Eds.), *Neighborhood poverty: Policy implications in studying neighborhoods*. Russell Sage Foundation New York, NY.
- Byrnes, H. F., Chen, M.-J., Miller, B. A., & Maguin, E. (2007). The relative importance of mothers' and youth's neighborhood perceptions for youth alcohol use and delinquency. *Journal of Youth and Adolescence*, 36(5), 649–659. <https://doi.org/10.1007/s10964-006-9154-2>
- Cappadocia, M. C., Desrocher, M., Pepler, D., & Schroeder, J. H. (2009). Contextualizing the neurobiology of conduct disorder in an emotion dysregulation framework. *Clinical Psychology Review*, 29(6), 506–518. <https://doi.org/10.1016/j.cpr.2009.06.001>
- Casey, B. J., Cannonier, T., Conley, M. I., Cohen, A. O., Barch, D. M., Heitzeg, M. M. . . . Dale, A. M. (2018). The adolescent brain cognitive development (ABCD) study: Imaging acquisition across 21 sites. *Developmental Cognitive Neuroscience*, 32, 43–54. <https://doi.org/10.1016/j.dcn.2018.03.001>
- Carlozzi, N. E., Beaumont, J. L., Tulsy, D. S., & Gershon, R. C. (2015). The NIH Toolbox pattern comparison processing speed test: Normative data. *Archives of Clinical Neuropsychology*, 30(5), 359–368. <https://doi.org/10.1093/arclin/acv031>
- Carlozzi, N. E., Tulsy, D. S., Chiaravalloti, N. D., Beaumont, J. L., Weintraub, S., Conway, K. . . . Gershon, R. C. (2014). NIH Toolbox cognitive battery (NIHTB-CB): The NIHTB pattern comparison processing speed test. *Journal of the International Neuropsychological Society*, 20(6), 630. <https://doi.org/10.1017/S1355617714000319>
- Carlozzi, N. E., Tulsy, D. S., Kail, R. V., & Beaumont, J. L. (2013). VI. NIH Toolbox cognition battery (CB): Measuring processing speed. *Monographs of the Society for Research in Child Development*, 78(4), 88–102. <https://doi.org/10.1111/mono.12036>
- Caspi, C. E., Kawachi, I., Subramanian, S. V., Adamkiewicz, G., & Sorensen, G. (2012). The relationship between diet and perceived and objective access to supermarkets among low-income housing residents. *Social Science & Medicine*, 75(7), 1254–1262. <https://doi.org/10.1016/j.socscimed.2012.05.014>
- Celeux, G., & Soromenho, G. (1996). An entropy criterion for assessing the number of clusters in a mixture model. *Journal of Classification*, 13(2), 195–212. <https://doi.org/10.1007/BF01246098>
- Chen, E., Langer, D. A., Raphaelson, Y. E., & Matthews, K. A. (2004). Socioeconomic status and health in adolescents: The role of stress interpretations. *Child Development*, 75(4), 1039–1052. <https://doi.org/10.1111/j.1467-8624.2004.00724.x>
- Cicchetti, D. (2010). Resilience under conditions of extreme stress: A multilevel perspective. *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)*, 9(3), 145–154. <https://doi.org/10.1002/j.2051-5545.2010.tb00297.x>
- Cicchetti, D., & Lynch, M. (1993). Toward an ecological/transactional model of community violence and child maltreatment: Consequences for children's development. *Psychiatry-interpersonal and Biological Processes*, 56(1), 96–118. <https://doi.org/10.1080/00332747.1993.11024624>
- Clark, D. B., Fisher, C. B., Bookheimer, S., Brown, S. A., Evans, J. H., Hopfer, C. . . . Yurgelun-Todd, D. (2018). Biomedical ethics and clinical oversight in multisite observational neuroimaging studies with children and adolescents: The ABCD experience. *Developmental Cognitive Neuroscience*, 32, 143–154. <https://doi.org/10.1016/j.dcn.2017.06.005>
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. Lawrence Erlbaum Associates. <https://doi.org/10.1234/12345678>
- Cohodes, E. M., Kitt, E. R., Baskin-Sommers, A., & Gee, D. G. (2020). Influences of early-life stress on frontolimbic circuitry: Harnessing a dimensional approach to elucidate the effects of heterogeneity in stress exposure. *Developmental Psychobiology*, 63(2), 153–172. <https://doi.org/10.1002/dev.21969>
- Cole, E. R. (2009). Intersectionality and research in psychology. *American Psychologist*, 64(3), 170. <https://doi.org/10.1037/a0014564>
- Coley, R. L., Sims, J., Dearing, E., & Spielvogel, B. (2018). Locating economic risks for adolescent mental and behavioral health: Poverty and affluence in

- families, neighborhoods, and schools. *Child Development*, 89(2), 360–369. <https://doi.org/10.1111/cdev.12771>
- Colich, N. L., Sheridan, M. A., Humphreys, K. L., Wade, M., Tibu, F., Nelson, C. A. . . . McLaughlin, K. A.** (2021). Heightened sensitivity to the caregiving environment during adolescence: Implications for recovery following early-life adversity. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 62(8), 937–948. <https://doi.org/10.1111/jcpp.13347>
- Compton, W. M., Dowling, G. J., & Garavan, H.** (2019). Ensuring the best use of data: The adolescent brain cognitive development study. *JAMA Pediatrics*, 137(9), 809–810. <https://doi.org/10.1001/jamapediatrics.2019.2081>
- Copeland, W. E., Goldston, D. B., & Costello, E. J.** (2017). Adult associations of childhood suicidal thoughts and behaviors: A prospective, longitudinal analysis. *Journal of the American Academy of Child and Adolescent Psychiatry*, 56(11), 958–965.e4. <https://doi.org/10.1016/j.jaac.2017.08.015>
- Crenshaw, K.** (1993). Demarginalizing the intersection of race and sex: A Black feminist critique of antidiscrimination doctrine, feminist theory and antiracist politics. In Weisbert D. K. (Eds.), *Feminist legal theory: Foundations*. Temple University Press Philadelphia: 383–395.
- Cuellar, J., Jones, D. J., & Sterrett, E.** (2015). Examining parenting in the neighborhood context: A review. *Journal of Child and Family Studies*, 24(1), 195–219. <https://doi.org/10.1007/s10826-013-9826-y>
- Danese, A., & McEwen, B. S.** (2012). Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiology & Behavior*, 106(1), 29–39. <https://doi.org/10.1016/j.physbeh.2011.08.019>
- Danese, A., & Widom, C. S.** (2020). Objective and subjective experiences of child maltreatment and their relationships with psychopathology. *Nature Human Behaviour*, 4(8), 811–818. <https://doi.org/10.1038/s41562-020-0880-3>
- Dawson, C. T., Wu, W., Fennie, K. P., Ibañez, G., Cano, M.Á., Pettit, J. W. . . . Trepka, M. J.** (2019). Perceived neighborhood social cohesion moderates the relationship between neighborhood structural disadvantage and adolescent depressive symptoms. *Health & Place*, 56, 88–98. <https://doi.org/10.1016/j.healthplace.2019.01.001>
- Denham, S. A., Mitchell-Copeland, J., Strandberg, K., Auerbach, S., & Blair, K.** (1997). Parental contributions to preschoolers' emotional competence: Direct and indirect effects. *Motivation and Emotion*, 21(1), 65–86. <https://doi.org/10.1023/A:1024426431247>
- Dick, A. S., Lopez, D. A., Watts, A. L., Heeringa, S., Reuter, C., Bartsch, H. . . . Stuart, E. A.** (2021). Meaningful associations in the adolescent brain cognitive development study. *NeuroImage*, 239, 118262. <https://doi.org/10.1016/j.neuroimage.2021.118262>
- Dikmen, S. S., Bauer, P. J., Weintraub, S., Mungas, D., Slotkin, J., Beaumont, J. L. . . . Heaton, R. K.** (2014). Measuring episodic memory across the life-span: NIH Toolbox picture sequence memory test. *Journal of the International Neuropsychological Society*, 20(6), 611. <https://doi.org/10.1017/S1355617714000460>
- Dube, S. R., Fairweather, D., Pearson, W. S., Felitti, V. J., Anda, R. F., & Croft, J. B.** (2009). Cumulative childhood stress and autoimmune diseases in adults. *Psychosomatic Medicine*, 71(2), 243–250. <https://doi.org/10.1097/PSY.0b013e3181907888>
- Eccles, J. S., & Roeser, R. W.** (2009). Schools, academic motivation, and stage-environment fit. In Lerner R. M., & Steinberg L. (Eds.), *Handbook of adolescent psychology: Individual bases of adolescent development*. John Wiley & Sons Inc: 404–434. <https://doi.org/10.1002/9780470479193.adlpsy001013>
- Echeverria, S. E., Diez-Roux, A. V., & Link, B. G.** (2004). Reliability of self-reported neighborhood characteristics. *Journal of Urban Health*, 81(4), 682–701. <https://doi.org/10.1093/jurban/jth151>
- El-Sheikh, M., & Harger, J.** (2001). Appraisals of marital conflict and children's adjustment, health, and physiological reactivity. *Developmental Psychology*, 37(6), 875–885. <https://doi.org/10.1037/0012-1649.37.6.875>
- Evans, G. W., Li, D., & Whipple, S. S.** (2013). Cumulative risk and child development. *Psychological Bulletin*, 139(6), 1342–1396. <https://doi.org/10.1037/a0031808>
- Fairchild, G., Hawes, D. J., Frick, P. J., Copeland, W. E., Odgers, C. L., Franke, B. . . . De Brito, S. A.** (2019). Conduct disorder. *Nature Reviews Disease Primers*, 5(1), 1–25. <https://doi.org/10.1038/s41572-019-0095-y>
- Fan, J., McCandliss, B. D., Sommer, T., Raz, A., & Posner, M. I.** (2002). Testing the efficiency and independence of attentional networks. *Journal of Cognitive Neuroscience*, 14(3), 340–347. <https://doi.org/10.1162/089892902317361886>
- Fay-Stammach, T., Hawes, D. J., & Meredith, P.** (2014). Parenting influences on executive function in early childhood: A review. *Child Development Perspectives*, 8(4), 258–264. <https://doi.org/10.1111/cdep.12095>
- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V. . . . Marks, J. S.** (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14(4), 245–258. [https://doi.org/10.1016/s0749-3797\(98\)00017-8](https://doi.org/10.1016/s0749-3797(98)00017-8)
- Fergusson, D. M., Horwood, L. J., & Ridder, E. M.** (2005). Show me the child at seven: The consequences of conduct problems in childhood for psychosocial functioning in adulthood. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 46(8), 837–849. <https://doi.org/10.1111/j.1469-7610.2004.00387.x>
- Fite, P. J., Vitulano, M., Wynn, P., Wimsatt, A., Gaertner, A., & Rathert, J.** (2010). Influence of perceived neighborhood safety on proactive and reactive aggression. *Journal of Community Psychology*, 38(6), 757–768. <https://doi.org/10.1002/jcop.20393>
- Friese, B., Grube, J. W., & Seninger, S.** (2015). Drinking among Native American and White Youths: The role of perceived neighborhood and school environment. *Journal of Ethnicity in Substance Abuse*, 14(3), 287–307. <https://doi.org/10.1080/15332640.2014.994723>
- Friedman, M., & Sharkey, P.** (2015). Violence and neighborhood disadvantage after the crime decline. *The Annals of the American Academy of Political and Social Science*, 660(1), 341–358. <https://doi.org/10.1177/0002716215579825>
- Furstenberg, F. F., Cook, T. D., Eccles, J., & Elder, G. H.** (2000). *Managing to make it: Urban families and adolescent success*. University of Chicago Press.
- Garavan, H., Bartsch, H., Conway, K., Decastro, A., Goldstein, R. Z., Heeringa, S. . . . Zahs, D.** (2018). Recruiting the ABCD sample: Design considerations and procedures. *Developmental Cognitive Neuroscience*, 32, 16–22. <https://doi.org/10.1016/j.dcn.2018.04.004>
- Gard, A. M., Waller, R., Shaw, D. S., Forbes, E. E., Hariri, A. R., & Hyde, L. W.** (2017). The long reach of early adversity: Parenting, stress, and neural pathways to antisocial behavior in adulthood. *Biological Psychiatry. Cognitive Neuroscience and Neuroimaging*, 2(7), 582–590. <https://doi.org/10.1016/j.bpsc.2017.06.005>
- Garnezy, N.** (1991). Resiliency and vulnerability to adverse developmental outcomes associated with poverty. *The American Behavioral Scientist*, 34(4), 416–430. <https://doi.org/10.1177/0002764291034004003>
- Gershon, R. C., Cook, K. F., Mungas, D., Manly, J. J., Slotkin, J., Beaumont, J. L. . . . Weintraub, S.** (2014). Language measures of the NIH Toolbox cognition battery. *Journal of the International Neuropsychological Society*, 20(6), 642–651. <https://doi.org/10.1017/S1355617714000411>
- Gershon, R. C., Slotkin, J., Manly, J. J., Blitz, D. L., Beaumont, J. L., Schnipke, D. . . . Weintraub, S.** (2013). IV. NIH Toolbox cognition battery (CB): Measuring language (vocabulary comprehension and reading decoding). *Monographs of the Society for Research in Child Development*, 78(4), 49–69. <https://doi.org/10.1111/mono.12034>
- Germiné, L., Robinson, E. B., Smoller, J. W., Calkins, M. E., Moore, T. M., Hakonarson, H. . . . Gur, R. E.** (2016). Association between polygenic risk for schizophrenia, neurocognition and social cognition across development. *Translational Psychiatry*, 6(10), e924. <https://doi.org/10.1038/tp.2016.147>
- Goldman-Mellor, S., Margerison-Zilko, C., Allen, K., & Cerda, M.** (2016). Perceived and objectively-measured neighborhood violence and adolescent psychological distress. *Journal of Urban Health: Bulletin of the New York Academy of Medicine*, 93(5), 758–769. <https://doi.org/10.1007/s11524-016-0079-0>
- Goodman, R., Meltzer, H., & Bailey, V.** (1998). The Strengths and Difficulties Questionnaire: A pilot study on the validity of the self-report version. *European Child & Adolescent Psychiatry*, 7(3), 125–130. <https://doi.org/10.1007/s007870050057>
- Goodman, R., & Scott, S.** (1999). Comparing the Strengths and Difficulties Questionnaire and the Child Behavior Checklist: Is small beautiful? *Journal of Abnormal Child Psychology*, 27(1), 17–24. <https://doi.org/10.1023/a:1022658222914>

- Gotlib, I. H., & Joormann, J.** (2010). Cognition and depression: Current status and future directions. *Annual Review of Clinical Psychology*, 6, 285–312. <https://doi.org/10.1146/annurev.clinpsy.121208.131305>
- Green, J. G., McLaughlin, K. A., Berglund, P. A., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., . . . Kessler, R. C.** (2010). Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication I: Associations with first onset of DSM-IV disorders. *Archives of General Psychiatry*, 67(2), 113–123. <https://doi.org/10.1001/archgenpsychiatry.2009.186>
- Greene, R. W., Biederman, J., Zerwas, S., Monuteaux, M. C., Goring, J. C., & Faraone, S. V.** (2002). Psychiatric comorbidity, family dysfunction, and social impairment in referred youth with oppositional defiant disorder. *The American Journal of Psychiatry*, 159(7), 1214–1224. <https://doi.org/10.1176/appi.ajp.159.7.1214>
- Gunnar, M., & Quevedo, K.** (2007). The neurobiology of stress and development. *Annual Review of Psychology*, 58, 145–173. <https://doi.org/10.1146/annurev.psych.58.110405.085605>
- Guyer, A. E., Kaufman, J., Hodgdon, H. B., Masten, C. L., Jazbec, S., Pine, D. S., . . . Ernst, M.** (2006). Behavioral alterations in reward system function: The role of childhood maltreatment and psychopathology. *Journal of the American Academy of Child & Adolescent Psychiatry*, 45(9), 1059–1067. <https://doi.org/10.1097/01.chi.0000227882.50404.11>
- Hadley-Ives, E., Stiffman, A. R., Elze, D., Johnson, S. D., & Dore, P.** (2000). Measuring neighborhood and school environments perceptual and aggregate approaches. *Journal of Human Behavior in the Social Environment*, 3(1), 1–28. https://doi.org/10.1300/J137v03n01_01
- Hale, L., Hill, T. D., Friedman, E., Nieto, F. J., Galvao, L. W., Engelman, C. D., . . . Peppard, P. E.** (2013). Perceived neighborhood quality, sleep quality, and health status: Evidence from the Survey of the Health of Wisconsin. *Social Science & Medicine*, 79, 16–22. <https://doi.org/10.1016/j.socscimed.2012.07.021>
- Harkness, K. L., Bruce, A. E., & Lumley, M. N.** (2006). The role of childhood abuse and neglect in the sensitization to stressful life events in adolescent depression. *Journal of Abnormal Psychology*, 115(4), 730–741. <https://doi.org/10.1037/0021-843X.115.4.730>
- Hasler, B. P., Kirisci, L., & Clark, D. B.** (2016). Restless sleep and variable sleep timing during late childhood accelerate the onset of alcohol and other drug involvement. *Journal of Studies on Alcohol and Drugs*, 77(4), 649–655. <https://doi.org/10.15288/jsad.2016.77.649>
- Herba, C. M., Ferdinand, R. F., van der Ende, J., & Verhulst, F. C.** (2007). Long-term associations of childhood suicide ideation. *Journal of the American Academy of Child and Adolescent Psychiatry*, 46(11), 1473–1481. <https://doi.org/10.1097/chi.0b013e318149e66f>
- Herrenkohl, T. I., & Herrenkohl, R. C.** (2007). Examining the overlap and prediction of multiple forms of child maltreatment, stressors, and socioeconomic status: A longitudinal analysis of youth outcomes. *Journal of Family Violence*, 22(7), 553–562. <https://doi.org/10.1007/s10896-007-9107-x>
- Hofferth, S. L., & Sandberg, J. F.** (2001). How American children spend their time. *Journal of Marriage and the Family*, 63(2), 295–308. <https://doi.org/10.1111/j.1741-3737.2001.00295.x>
- Hoffman, E. A., Clark, D. B., Orendain, N., Hudziak, J., Squeglia, L. M., & Dowling, G. J.** (2019). Stress exposures, neurodevelopment and health measures in the ABCD study. *Neurobiology of Stress*, 10, 100157. <https://doi.org/10.1016/j.ynstr.2019.100157>
- Holmes, C. J., Kim-Spoon, J., & Deater-Deckard, K.** (2016). Linking executive function and peer problems from early childhood through middle adolescence. *Journal of Abnormal Child Psychology*, 44(1), 31–42. <https://doi.org/10.1007/s10802-015-0044-5>
- Hostinar, C. E., & Miller, G. E.** (2019). Protective factors for youth confronting economic hardship: Current challenges and future avenues in resilience research. *The American Psychologist*, 74(6), 641–652. <https://doi.org/10.1037/amp0000520>
- Howard, M. C., & Hoffman, M. E.** (2018). Variable-centered, person-centered, and person-specific approaches: where theory meets the method. *Organizational Research Methods*, 21(4), 846–876. <https://doi.org/10.1177/1094428117744021>
- Humphreys, K. L., Katz, S. J., Lee, S. S., Hammen, C., Brennan, P. A., & Najman, J. M.** (2013). The association of ADHD and depression: Mediation by peer problems and parent-child difficulties in two complementary samples. *Journal of Abnormal Psychology*, 122(3), 854–867. <https://doi.org/10.1037/a0033895>
- Iacono, W. G., Heath, A. C., Hewitt, J. K., Neale, M. C., Banich, M. T., Luciana, M. M., . . . Bjork, J. M.** (2018). The utility of twins in developmental cognitive neuroscience research: How twins strengthen the ABCD research design. *Developmental Cognitive Neuroscience*, 32, 30–42. <https://doi.org/10.1016/j.dcn.2017.09.001>
- Jackson, C. L., Redline, S., & Emmons, K. M.** (2015). Sleep as a potential fundamental contributor to disparities in cardiovascular health. *Annual Review of Public Health*, 36, 417–440. <https://doi.org/10.1146/annurev-publhealth-031914-122838>
- Jamieson, J. P., Hangen, E. J., Lee, H. Y., & Yeager, D. S.** (2018). Capitalizing on appraisal processes to improve affective responses to social stress. *Emotion Review: Journal of the International Society for Research on Emotion*, 10(1), 30–39. <https://doi.org/10.1177/1754073917693085>
- Jernigan, T. L., Brown, S. A., & ABCD Consortium Coordinators** (2018). Introduction. *Developmental Cognitive Neuroscience*, 32, 1–3. <https://doi.org/10.1016/j.dcn.2018.02.002>
- Kasen, S., Johnson, J., & Cohen, P.** (1990). The impact of school emotional climate on student psychopathology. *Journal of Abnormal Child Psychology*, 18(2), 165–177. <https://doi.org/10.1007/BF00910728>
- Kaufman, J., Birmaher, B., Axelson, D., Perepletchikova, F., Brent, D., & Ryan, N.** Kiddie Schedule for Affective Disorders and Schizophrenia Present and Lifetime Version 2013: Working Draft (KSADS-PL). 2013.
- Kaufman, J., Birmaher, B., Brent, D., Rao, U., Flynn, C., Moreci, P., . . . Ryan, N.** (1997). Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (K-SADS-PL): Initial reliability and validity data. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(7), 980–988. <https://doi.org/10.1097/00004583-199707000-00021>
- Kawabata, Y., Alink, L. R. A., Tseng, W.-L., van IJzendoorn, M. H., & Crick, N. R.** (2011). Maternal and paternal parenting styles associated with relational aggression in children and adolescents: A conceptual analysis and meta-analytic review. *Developmental Review*, 31(4), 240–278. <https://doi.org/10.1016/j.dr.2011.08.001>
- Klimes-Dougan, B., & Garber, J.** (2016). Regulatory control and depression in adolescents: Findings from neuroimaging and neuropsychological research. *Journal of Clinical Child and Adolescent Psychology: The Official Journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division*, 45(1), 1–5. <https://doi.org/10.1080/15374416.2015.1123637>
- Knoll, L. J., Magis-Weinberg, L., Speekenbrink, M., & Blakemore, S.-J.** (2015). Social influence on risk perception during adolescence. *Psychological Science*, 26(5), 583–592. <https://doi.org/10.1177/0956797615569578>
- Kobak, K. A., Kratochvil, C. J., Stanger, C., & Kaufman, J.** (2013). Computerized screening of comorbidity in adolescents with substance or psychiatric disorders. *Anxiety Disorders and Depression (La Jolla, CA)*.
- Kohen, D. E., Brooks-Gunn, J., Leventhal, T., & Hertzman, C.** (2002). Neighborhood income and physical and social disorder in Canada: Associations with young children's competencies. *Child Development*, 73(6), 1844–1860. <https://doi.org/10.1111/1467-8624.t01-1-00510>
- Kuhlman, K. R., Chiang, J. J., Horn, S., & Bower, J. E.** (2017). Developmental psychoneuroendocrine and psychoneuroimmune pathways from childhood adversity to disease. *Neuroscience and Biobehavioral Reviews*, 80, 166–184. <https://doi.org/10.1016/j.neubiorev.2017.05.020>
- Lambert, S. F., Brown, T. L., Phillips, C. M., & Ialongo, N. S.** (2004). The relationship between perceptions of neighborhood characteristics and substance use among urban African American adolescents. *American Journal of Community Psychology*, 34(3-4), 205–218. <https://doi.org/10.1007/s10464-004-7415-3>
- LaRusso, M. D., Romer, D., & Selman, R. L.** (2007). Teachers as builders of respectful school climates: Implications for adolescent drug use norms and depressive symptoms in high school. *Journal of Youth and Adolescence*, 37(4), 386. <https://doi.org/10.1007/s10964-007-9212-4>
- Latham, R. M., Mark, K. M., & Oliver, B. R.** (2017). A harsh parenting team? Maternal reports of coparenting and coercive parenting interact in association with children's disruptive behaviour. *Journal of Child Psychology and*

- Psychiatry, and Allied Disciplines*, 58(5), 603–611. <https://doi.org/10.1111/jcpp.12665>
- Lenth, R. V.** Estimated Marginal Means, aka Least-Squares Means [R package emmeans version 1.7.0]. 2021, <https://CRAN.R-project.org/package=emmeans>
- Lerner, R. M.** (1991). Changing organism–context relations as the basic process of development: A developmental contextual perspective. *Developmental Psychology*, 27(1), 27–32. <https://doi.org/10.1037/0012-1649.27.1.27>
- Lezak, M. D., Howieson, D. B., Loring, D. W., & Fischer, J. S.** (2004). *Neuropsychological assessment*. Oxford University Press.
- Li, A., & Fischer, M. J.** (2017). Advantaged/disadvantaged school neighborhoods, parental networks, and parental involvement at elementary school. *Sociology of Education*, 90(4), 355–377. <https://doi.org/10.1177/0038040717732332>
- Luciana, M., Bjork, J. M., Nagel, B. J., Barch, D. M., Gonzalez, R., Nixon, S. J., . . . Banich, M. T.** (2018). Adolescent neurocognitive development and impacts of substance use: Overview of the adolescent brain cognitive development (ABCD) baseline neurocognition battery. *Developmental Cognitive Neuroscience*, 32, 67–79. <https://doi.org/10.1016/j.dcn.2018.02.006>
- Lüdtke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D.** (2021). Performance: An R package for assessment, comparison and testing of statistical models. *Journal of Open Source Software*, 6(60), 3139. <https://doi.org/10.21105/joss.03139>
- Lupien, S. J., McEwen, B. S., Gunnar, M. R., & Heim, C.** (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews: Neuroscience*, 10(6), 434–445. <https://doi.org/10.1038/nrn2639>
- Magnusson, D., & Stattin, H.** (1998). Person-context interaction theories. In Damon W., (Eds.), *Handbook of child psychology: Theoretical models of human development*. vol. 1, John Wiley & Sons Inc: 685–759.
- Manly, J. T., Kim, J. E., Rogosch, F. A., & Cicchetti, D.** (2001). Dimensions of child maltreatment and children’s adjustment: Contributions of developmental timing and subtype. *Development and Psychopathology*, 13(4), 759–782.
- Mason, M., Cheung, I., & Walker, L.** (2004). Substance use, social networks, and the geography of urban adolescents. *Substance Use & Misuse*, 39(10–12), 1751–1777.
- Masi, G., Milone, A., Pisano, S., Lenzi, F., Muratori, P., Gemo, I. . . Vicari, S.** (2014). Emotional reactivity in referred youth with disruptive behavior disorders: The role of the callous-unemotional traits. *Psychiatry Research*, 220(1–2), 426–432. <https://doi.org/10.1016/j.psychres.2014.07.035>
- Masten, A. S., & Cicchetti, D.** (2010). Developmental cascades. *Development and Psychopathology*, 22(3), 491–495. <https://doi.org/10.1017/S0954579410000222>
- Masten, A. S.** (2001). Ordinary magic. Resilience processes in development. *The American Psychologist*, 56(3), 227–238. <https://doi.org/10.1037/0003-066x.56.3.227>
- Masten, A. S., & Narayan, A. J.** (2012). Child development in the context of disaster, war, and terrorism: Pathways of risk and resilience. *Annual Review of Psychology*, 63, 227–257. <https://doi.org/10.1146/annurev-psych-120710-100356>
- Masyn, K. E.** (2013). Latent class analysis and finite mixture modeling. In Little T. D. (Eds.), *The Oxford handbook of quantitative methods in psychology: Vol. 2: Statistical analysis*. Oxford University Press, Inc, <https://doi.org/10.1093/oxfordhb/9780199934898.013.0025>
- Matthys, W., Vanderschuren, L. J. M. J., Schutter, D. J. L. G., & Lochman, J. E.** (2012). Impaired neurocognitive functions affect social learning processes in oppositional defiant disorder and conduct disorder: Implications for interventions. *Clinical Child and Family Psychology Review*, 15(3), 234–246. <https://doi.org/10.1007/s10567-012-0118-7>
- Maughan, B.** (1995). Annotation: Long-term outcomes of developmental reading problems. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 36(3), 357–371. <https://doi.org/10.1111/j.1469-7610.1995.tb01296.x>
- Mayne, S. L., Mitchell, J. A., Virudachalam, S., Fiks, A. G., & Williamson, A. A.** (2021). Neighborhood environments and sleep among children and adolescents: A systematic review. *Sleep Medicine Reviews*, 57, 101465. <https://doi.org/10.1016/j.smrv.2021.101465>
- McCoy, D. C., Roy, A. L., & Raver, C. C.** (2016). Neighborhood crime as a predictor of individual differences in emotional processing and regulation. *Developmental Science*, 19(1), 164–174. <https://doi.org/10.1111/desc.12287>
- McEwen, B. S., & Wingfield, J. C.** (2003). The concept of allostasis in biology and biomedicine. *Hormones and Behavior*, 43(1), 2–15. [https://doi.org/10.1016/s0018-506x\(02\)00024-7](https://doi.org/10.1016/s0018-506x(02)00024-7)
- McLaughlin, K. A., & Lambert, H. K.** (2017). Child trauma exposure and psychopathology: Mechanisms of risk and resilience. *Current Opinion in Psychology*, 14, 29–34. <https://doi.org/10.1016/j.copsyc.2016.10.004>
- McNulty, T. L., & Bellair, P. E.** (2003). Explaining racial and ethnic differences in serious adolescent violent behavior. *Criminology: An Interdisciplinary Journal*, 41(3), 709–747. <https://doi.org/10.1111/j.1745-9125.2003.tb01002.x>
- Meldrum, R. C., Jackson, D. B., Archer, R., & Ammons-Blanford, C.** (2018). Perceived school safety, perceived neighborhood safety, and insufficient sleep among adolescents. *Sleep Health*, 4(5), 429–435. <https://doi.org/10.1016/j.sleh.2018.07.006>
- Millam, A. J., Furr-Holden, C. D. M., & Leaf, P. J.** (2010). Perceived school and neighborhood safety, neighborhood violence and academic achievement in urban school children. *The Urban Review*, 42(5), 458–467. <https://doi.org/10.1007/s11256-010-0165-7>
- Minh, A., Muhajarine, N., Janus, M., Brownell, M., & Guhn, M.** (2017). A review of neighborhood effects and early child development: How, where, and for whom, do neighborhoods matter? *Health & Place*, 46, 155–174. <https://doi.org/10.1016/j.healthplace.2017.04.012>
- Mogg, K., & Bradley, B. P.** (2005). Attentional bias in generalized anxiety disorder versus depressive disorder. *Cognitive Therapy and Research*, 29(1), 29–45. <https://doi.org/10.1007/s10608-005-1646-y>
- Moos, R. H., & Moos, B. S.** (1994). *Family environment scale manual* (3rd edn. Consulting Psychologists Press, Palo Alto, CA.
- Morris, A. S., Silk, J. S., Steinberg, L., Myers, S. S., & Robinson, L. R.** (2007). The role of the family context in the development of emotion regulation. *Social Development*, 16(2), 361–388. <https://doi.org/10.1111/j.1467-9507.2007.00389.x>
- Mrug, S., & Windle, M.** (2010). Prospective effects of violence exposure across multiple contexts on early adolescents’ internalizing and externalizing problems. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 51(8), 953–961. <https://doi.org/10.1111/j.1469-7610.2010.02222.x>
- Mujahid, M. S., Diez Roux, A. V., Morenoff, J. D., & Raghunathan, T.** (2007). Assessing the measurement properties of neighborhood scales: From psychometrics to econometrics. *American Journal of Epidemiology*, 165(8), 858–867. <https://doi.org/10.1093/aje/kwm040>
- Nails, A. M., Mullis, R. L., & Mullis, A. K.** (2009). American Indian youth’s perceptions of their environment and their reports of depressive symptoms and alcohol/marijuana use. *Adolescence*, 44(176), 965–978.
- Ogilvie, J. M., Stewart, A. L., Chan, R. C. K., & Shum, D. H. K.** (2011). Neuropsychological measures of executive function and antisocial behavior: A meta-analysis: Executive function and antisocial behavior. *Criminology: An Interdisciplinary Journal*, 49(4), 1063–1107. <https://doi.org/10.1111/j.1745-9125.2011.00252.x>
- Oliver, B. R.** (2015). Unpacking externalising problems: Negative parenting associations for conduct problems and irritability. *BJPsych Open*, 1(1), 42–47. <https://doi.org/10.1192/bjpo.bp.115.000125>
- Orstad, S. L., McDonough, M. H., Stapleton, S., Altincekic, C., & Troped, P. J.** (2017). A systematic review of agreement between perceived and objective neighborhood environment measures and associations with physical activity outcomes. *Environment and Behavior*, 49(8), 904–932. <https://doi.org/10.1177/0013916516670982>
- Orth, U., Clark, D. A., Donnellan, M. B., & Robins, R. W.** (2021). Testing prospective effects in longitudinal research: Comparing seven competing cross-lagged models. *Journal of Personality and Social Psychology*, 120(4), 1013–1034. <https://doi.org/10.1037/pspp0000358>
- Owens, M. M., Potter, A., Hyatt, C. S., Albaugh, M., Thompson, W. K., Jernigan, T. . . . Garavan, H.** (2021). Recalibrating expectations about effect size: A multi-method survey of effect sizes in the ABCD study. *PLoS One*, 16(9), e0257535. <https://doi.org/10.1371/journal.pone.0257535>
- Padgaonkar, N. T., Baker, A. E., Dapretto, M., Galván, A., Frick, P. J., Steinberg, L. . . . Cauffman, E.** (2021). Exploring disproportionate minority

- contact in the juvenile justice system over the year following first arrest. *Journal of Research on Adolescence: The Official Journal of the Society for Research on Adolescence*, 31(2), 317–334. <https://doi.org/10.1111/jora.12599>
- Palacios-Barrios, E. E., & Hanson, J. L. (2019). Poverty and self-regulation: Connecting psychosocial processes, neurobiology, and the risk for psychopathology. *Comprehensive Psychiatry*, 90, 52–64. <https://doi.org/10.1016/j.comppsy.2018.12.012>
- Parke, R. D. (1994). Progress, paradigms, and unresolved problems: A commentary on recent advances in our understanding of children's emotions. *Merrill-Palmer Quarterly*, 40(1), 157–169.
- Patterson, G. R. (1982). *Coercive family process*, vol. 3. Castalia Publishing Company.
- Pinquart, M. (2017). Associations of parenting dimensions and styles with externalizing problems of children and adolescents: An updated meta-analysis. *Developmental Psychology*, 53(5), 873–932. <https://doi.org/10.1037/dev0000295>
- Pollak, S. (2015). Multilevel developmental approaches to understanding the effects of child maltreatment: Recent advances and future challenges. *Development and Psychopathology*, 27(4pt2), 1387–1397. <https://doi.org/10.1017/S0954579415000826>
- Puliafico, A. C., & Kendall, P. C. (2006). Threat-related attentional bias in anxious youth: A review. *Clinical Child and Family Psychology Review*, 9(3), 162–180. <https://doi.org/10.1007/s10567-006-0009-x>
- R Core Team (2020). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>
- Rasmussen, A., Aber, M. S., & Bhana, A. (2004). Adolescent coping and neighborhood violence: Perceptions, exposure, and urban youth's efforts to deal with danger. *American Journal of Community Psychology*, 33(1-2), 61–75. <https://doi.org/10.1023/b:ajcp.0000014319.32655.66>
- Repetti, R. L., Taylor, S. E., & Seeman, T. E. (2002). Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin*, 128(2), 330–366. <https://doi.org/10.1037//0033-2909.128.2.330>
- Rivenbark, J. G., Odgers, C. L., Caspi, A., Harrington, H., Hogan, S., Houts, R. M., . . . Moffitt, T. E. (2018). The high societal costs of childhood conduct problems: Evidence from administrative records up to age 38 in a longitudinal birth cohort. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 59(6), 703–710. <https://doi.org/10.1111/jcpp.12850>
- Roberts, R. E., & Duong, H. T. (2017). Is there an association between short sleep duration and adolescent anxiety disorders? *Sleep Medicine*, 30, 82–87. <https://doi.org/10.1016/j.sleep.2016.02.007>
- Romano, E., Tremblay, R. E., Boulerice, B., & Swisher, R. (2005). Multilevel correlates of childhood physical aggression and prosocial behavior. *Journal of Abnormal Child Psychology*, 33(5), 565–578. <https://doi.org/10.1007/s10802-005-6738-3>
- Roosa, M. W., Jones, S., Tein, J.-Y., & Cree, W. (2003). Prevention science and neighborhood influences on low-income children's development: Theoretical and methodological issues. *American Journal of Community Psychology*, 31(1-2), 55–72. <https://doi.org/10.1023/a:1023070519597>
- Roosa, M. W., White, R. M. B., Zaiders, K. H., & Tein, J.-Y. (2009). An examination of the role of perceptions in neighborhood research. *Journal of Community Psychology*, 37(3), 327–341. <https://doi.org/10.1002/jcop.20298>
- Rosenberg, J., Beymer, P., Anderson, D., van Lissa, C. J., & Schmidt, J. (2018). TidyLPA: An R package to easily carry out latent profile analysis (LPA) using open-source or commercial software. *Journal of Open Source Software*, 3(30), 978. <https://doi.org/10.21105/joss.00978>
- Rosenfield, D., Jouriles, E. N., McDonald, R., & Mueller, V. (2014). Interparental conflict, community violence, and child problems: Making sense of counterintuitive findings. *The American Journal of Orthopsychiatry*, 84(3), 275–283. <https://doi.org/10.1037/h0099805>
- Rudolph, K. D., Monti, J. D., Modi, H., Sze, W. Y., & Troop-Gordon, W. (2020). Protecting youth against the adverse effects of peer victimization: Why do parents matter? *Journal of Abnormal Child Psychology*, 48(2), 163–176. <https://doi.org/10.1007/s10802-019-00576-9>
- Sampson, R. J., Sharkey, P., & Raudenbush, S. W. (2008). Durable effects of concentrated disadvantage on verbal ability among African-American children. *Proceedings of the National Academy of Sciences of the United States of America*, 105(3), 845–852. <https://doi.org/10.1073/pnas.0710189104>
- Schielzeth, H., Dingemans, N. J., Nakagawa, S., Westneat, D. F., Allogue, H., Teplitsky, C., . . . Araya-Ajoy, Y. G. (2020). Robustness of linear mixed-effects models to violations of distributional assumptions. *Methods in Ecology and Evolution*, 11(9), 1141–1152. <https://doi.org/10.1111/2041-210X.13434>
- Schwarz, G. (1978). Estimating the dimension of a model. *The Annals of Statistics*, 6(2), 461–464.
- Scrucca, L., Fop, M., Murphy, T. B., & Raftery, A. E. (2016). mclust 5: Clustering, classification and density estimation using Gaussian finite mixture models. *The R Journal*, 8(1), 289–317.
- Shackman, J. E., & Pollak, S. D. (2014). Impact of physical maltreatment on the regulation of negative affect and aggression. *Development and Psychopathology*, 26(4pt1), 1021–1033. <https://doi.org/10.1017/S0954579414000546>
- Shan, Z., Ma, H., Xie, M., Yan, P., Guo, Y., Bao, W., . . . Liu, L. (2015). Sleep duration and risk of type 2 diabetes: A meta-analysis of prospective studies. *Diabetes Care*, 38(3), 529–537. <https://doi.org/10.2337/dc14-2073>
- Sharkey, P. (2010). The acute effect of local homicides on children's cognitive performance. *Proceedings of the National Academy of Sciences of the United States of America*, 107(26), 11733–11738. <https://doi.org/10.1073/pnas.1000690107>
- Sheridan, M. A., & McLaughlin, K. A. (2014). Dimensions of early experience and neural development: Deprivation and threat. *Trends in Cognitive Sciences*, 18(11), 580–585. <https://doi.org/10.1016/j.tics.2014.09.001>
- Silk, J. S., Sessa, F. M., Morris, A. S., Steinberg, L., & Avenevoli, S. (2004). Neighborhood cohesion as a buffer against hostile maternal parenting. *Journal of Family Psychology: JFP: Journal of the Division of Family Psychology of the American Psychological Association*, 18(1), 135–146. <https://doi.org/10.1037/0893-3200.18.1.135>
- Simmons, C., Conley, M. I., Gee, D. G., Baskin-Sommers, A., Barch, D. M., Hoffman, E. A., . . . Casey, B. J. (2021). Responsible use of open-access developmental data: The Adolescent Brain Cognitive Development (ABCD) study. *Psychological Science*, 32(6), 866–870. <https://doi.org/10.1177/09567976211003564>
- Smith, K. E., & Pollak, S. D. (2021). Rethinking concepts and categories for understanding the neurodevelopmental effects of childhood adversity. *Perspectives on Psychological Science: A Journal of the Association for Psychological Science*, 16(1), 67–93. <https://doi.org/10.1177/1745691620920725>
- St Clair, M. C., Croudace, T., Dunn, V. J., Jones, P. B., Herbert, J., & Goodyer, I. M. (2015). Childhood adversity subtypes and depressive symptoms in early and late adolescence. *Development and Psychopathology*, 27(3), 885–899. <https://doi.org/10.1017/S0954579414000625>
- Steinberg, L. (2005). Cognitive and affective development in adolescence. *Trends in Cognitive Sciences*, 9(2), 69–74. <https://doi.org/10.1016/j.tics.2004.12.005>
- Tandon, M., Cardeli, E., & Luby, J. (2009). Internalizing disorders in early childhood: A review of depressive and anxiety disorders. *Child and Adolescent Psychiatric Clinics of North America*, 18(3), 593–610. <https://doi.org/10.1016/j.chc.2009.03.004>
- Telzer, E. H., Fuligni, A. J., Lieberman, M. D., & Galván, A. (2013). The effects of poor quality sleep on brain function and risk taking in adolescence. *NeuroImage*, 71, 275–283. <https://doi.org/10.1016/j.neuroimage.2013.01.025>
- Thapa, A., Cohen, J., Guffey, S., & Higgins-D'Alessandro, A. (2013). A review of school climate research. *Review of Educational Research*, 83(3), 357–385.
- Usami, S., Todo, N., & Murayama, K. (2019). Modeling reciprocal effects in medical research: Critical discussion on the current practices and potential alternative models. *PLoS One*, 14(9), e0209133. <https://doi.org/10.1371/journal.pone.0209133>
- Verbeke, G., & Molenberghs, G. (2013). *Linear mixed models for longitudinal data*. Springer, New York.
- Vermunt, J. K. (2010). Latent class modeling with covariates: Two improved three-step approaches. *Political Analysis*, 18(4), 450–469. <https://doi.org/10.1093/pan/mpq025>

- Vermunt, J. K., & Magidson, J. (2002). Latent class cluster analysis. In Hagenaars J. A., & McCutcheon A. L. (Eds.), *Applied latent class analysis*. Cambridge University Press: 89–106.
- Voisin, D. R., Hartly, J., Kim, D. H., Elsaesser, C., & Takahashi, L. M. (2017). Assessing the relationship between parental influences and wellbeing among low income African American adolescents in Chicago. *Child & Youth Care Forum*, 46(2), 223–242. <https://doi.org/10.1007/s10566-016-9373-y>
- Wade, M., Zeanah, C. H., Fox, N. A., Tibu, F., Ciolan, L. E., & Nelson, C. A. (2019). Stress sensitization among severely neglected children and protection by social enrichment. *Nature Communications*, 10(1), 5771. <https://doi.org/10.1038/s41467-019-13622-3>
- Wagner, S., Müller, C., Helmreich, I., Huss, M., & Tadić, A. (2015). A meta-analysis of cognitive functions in children and adolescents with major depressive disorder. *European Child & Adolescent Psychiatry*, 24(1), 5–19. <https://doi.org/10.1007/s00787-014-0559-2>
- Wang, M.-T., & Degol, J. L. (2016). School climate: A review of the construct, measurement, and impact on student outcomes. *Educational Psychology Review*, 28(2), 315–352. <https://doi.org/10.1007/s10648-015-9319-1>
- Weller, B. E., Bowen, N. K., & Faubert, S. J. (2020). Latent class analysis: A guide to best practice. *The Journal of Black Psychology*, 46(4), 287–311. <https://doi.org/10.1177/0095798420930932>
- Wigfield, A., Byrnes, J. P., & Eccles, J. S. (2006). Development during early and middle adolescence. In P. A. Alexander & P. H. Winne (Eds.), *Handbook of educational psychology* (pp. 87–113). Lawrence Erlbaum Associates Publishers.
- Wiggins, J. L., Mitchell, C., Hyde, L. W., & Monk, C. S. (2015). Identifying early pathways of risk and resilience: The co-development of internalizing and externalizing symptoms and the role of harsh parenting. *Development and Psychopathology*, 27(4 Pt 1), 1295. <https://doi.org/10.1017/S0954579414001412>
- Williams, D. R., & Collins, C. (2001). Racial residential segregation: A fundamental cause of racial disparities in health. *Public Health Reports (Washington, D.C.: 1974)*, 116(5), 404–416. <https://doi.org/10.1093/phr/116.5.404>
- Williams, David R., & Jackson, P. B. (2005). Social sources of racial disparities in health. *Health Affairs (Project Hope)*, 24(2), 325–334. <https://doi.org/10.1377/hlthaff.24.2.325>
- Williams, T. R., Davis, L. E., Cribbs, J. M., Saunders, J., & Williams, J. H. (2002). Friends, family, and neighborhood: Understanding academic outcomes of African American youth. *Urban Education*, 37(3), 408–431.
- Young, A. R., Beitchman, J. H., Johnson, C., Douglas, L., Atkinson, L., Escobar, M. . . . Wilson, B. (2002). Young adult academic outcomes in a longitudinal sample of early identified language impaired and control children. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 43(5), 635–645. <https://doi.org/10.1111/1469-7610.00052>
- Youngblade, L. M., Theokas, C., Schulenberg, J., Curry, L., Huang, I.-C., & Novak, M. (2007). Risk and promotive factors in families, schools, and communities: A contextual model of positive youth development in adolescence. *Pediatrics*, 119(Suppl 1), S47–S53. <https://doi.org/10.1542/peds.2006-2089H>
- Zucker, R. A., Gonzalez, R., Feldstein Ewing, S. W., Paulus, M. P., Arroyo, J., Fuligni, A. . . . Wills, T. (2018). Assessment of culture and environment in the Adolescent Brain and Cognitive Development Study: Rationale, description of measures, and early data. *Developmental Cognitive Neuroscience*, 32, 107–120. <https://doi.org/10.1016/j.dcn.2018.03.004>